

B7-Like Molecules and Uses Thereof

This application claims the benefit of U.S.

Provsional Application No. 60/214,512, filed June 28,

2000, which is hereby incorporated by reference

Field of the Invention

The present invention relates to novel B7-like polypeptides and nucleic acid molecules encoding the same. The invention also relates to vectors, host cells, pharmaceutical compositions, selective binding agents and methods for producing B7-like polypeptides. Also provided for are methods for the diagnosis, treatment, amelioration, and/or prevention of diseases associated with B7-like polypeptides.

15

20

25

30

10

Background of the Invention

Technical advances in the identification, cloning, expression and manipulation of nucleic acid molecules and the deciphering of the human genome have greatly accelerated the discovery of novel therapeutics. nucleic acid sequencing techniques can now generate sequence information at unprecedented rates coupled with computational analyses, allow the assembly overlapping sequences into partial and entire genomes and the identification of polypeptide-encoding regions. A comparison of a predicted amino acid sequence against a database compilation of known amino acid sequences allows one to determine the extent of homology to previously identified sequences and/or structural landmarks. The cloning and expression of a polypeptide-encoding region of a nucleic acid molecule provides a polypeptide product for structural and functional analyses. The manipulation of nucleic acid molecules and encoded polypeptides may confer

advantageous properties on a product for use as a therapeutic.

In spite of the significant technical advances in genome research over the past decade, the potential for the development of novel therapeutics based on the human genome is still largely unrealized. Many genes encoding potentially beneficial polypeptide therapeutics, or those encoding polypeptides, which may act as "targets" for therapeutic molecules, have still not been identified.

Accordingly, it is an object of the invention to identify novel polypeptides and nucleic acid molecules encoding the same, which have diagnostic or therapeutic benefit.

15

10

5

Summary of the Invention

The present invention relates to novel B7-like nucleic acid molecules and encoded polypeptides.

The invention provides for an isolated nucleic 20 acid molecule comprising a nucleotide sequence selected from the group consisting of:

- (a) the nucleotide sequence as set forth in SEQ ID NOs: 1, 3, 5 or 7;
- (b) the nucleotide sequence as set forth in SEQ 25 ID NOs: 9, 11 or 13;
 - (c) a nucleotide sequence encoding the polypeptide as set forth in SEQ ID NOs: 2, 4, 6 or 8;
 - (d) a nucleotide sequence encoding the polypeptide as set forth in SEQ ID NOs: 10, 12 or 14;
- 30 (e) a nucleotide sequence which hybridizes under

25

A-692

moderately or highly stringent conditions to the complement of (a) or (b), wherein the encoded polypeptide has an activity of the mature form of a polypeptide as set forth in SEQ ID NOs: 2, 4, 6 or 8;

- a nucleotide sequence which hybridizes under moderately orhighly stringent conditions complement of (a) or (b), wherein the encoded polypeptide has an activity of the mature form of a polypeptide as set forth in SEQ ID NOs: 10, 12 or 14; 10 and
 - (g) a nucleotide sequence complementary to any of (a)-(f).

The invention also provides for an isolated nucleic acid molecule comprising a nucleotide sequence selected from the group consisting of:

- (a) a nucleotide sequence encoding a polypeptide that is at least about 70, 75, 80, 85, 90, 95, 96, 97, 98, or 99 percent identical to the polypeptide as set forth in SEQ ID NOs: 2, 4, 6 or 8, wherein the polypeptide has an activity of the mature form of a polypeptide as set forth in SEQ ID NOs: 2, 4, 6 or 8;
- (b) a nucleotide sequence encoding a polypeptide that is at least about 70, 75, 80, 85, 90, 95, 96, 97, 98 or 99 percent identical to the polypeptide as set forth in SEQ ID NOs: 10, 12 or 14, wherein the polypeptide has an activity of the mature form of a polypeptide as set forth in SEQ ID NOs: 10, 12 or 14;
- (c) a nucleotide sequence encoding an allelic variant or splice variant of the nucleotide sequence as set forth in SEQ ID NOs: 1, 3, 5 or 7, wherein the

A-692

encoded polypeptide has an activity of the mature form of a polypeptide as set forth in SEQ ID NOs: 2, 4, 6 or 8;

- (d) a nucleotide sequence encoding an allelic variant or splice variant of the nucleotide sequence as set forth in SEQ ID NOs: 9, 11 or 13, wherein the encoded polypeptide has an activity of the mature form of a polypeptide as set forth in SEQ ID NOs: 10, 12 or 14;
- 10 (e) a nucleotide sequence of SEQ ID NOs: 1, 3, 5 or 7, or (a) or (b), above, encoding a polypeptide fragment of at least about 25 amino acid residues, wherein the polypeptide has an activity of the mature form of a polypeptide as set forth in SEQ ID NOs: 2, 4, 15 6 or 8;
- (f) a nucleotide sequence of SEQ ID NOs: 9, 11 or 13, or (a) or (b), above, encoding a polypeptide fragment of at least about 25 amino acid residues, wherein the polypeptide has an activity of the mature 20 form of a polypeptide as set forth in SEQ ID NOs: 10, 12 or 14;
 - g) a nucleotide sequence encoding a polypeptide that has a substitution and/or deletion of 1 to 100 amino acid residues as set forth in any of SEQ ID NOs: 9, 11 or 13, wherein the encoded polypeptide has an activity of the mature form of a polypeptide as set forth in SEQ ID NOs: 10, 12, or 14;
- h) a nucleotide sequence of SEQ ID NOs: 1, 3, 5
 or 7, or (a), (c), (e) or (f), above, comprising a
 fragment of at least about 16 nucleotides;
 - i) a nucleotide sequence of SEQ ID NOs: 9, 11 or

5

13, or (b), (d), (f) or (h), above, comprising a fragment of at least about 16 nucleotides;

- j) a nucleotide sequence which hybridizes under moderately or highly stringent conditions to the complement of any of (a), (c), (e), (g) or (i), above, wherein the polypeptide has an activity of the mature form of a polypeptide as set forth in SEQ ID NOs: 2, 4, 6 or 8;
- k) a nucleotide sequence which hybridizes under 10 moderately or highly stringent conditions to the complement of any of (b), (d), (f), (h) or (j), above, wherein the polypeptide has an activity of the mature form of a polypeptide as set forth in SEQ ID NOs: 10, 12 or 14; and
- 15 1) a nucleotide sequence complementary to any of (a)-(1).

The invention further provides for an isolated nucleic acid molecule comprising a nucleotide sequence selected from the group consisting of:

- 20 (a) a nucleotide sequence encoding a polypeptide as set forth in SEQ ID NOs: 2, 4, 6 or 8 with at least one conservative amino acid substitution, wherein the polypeptide has an activity of the mature form of a polypeptide as set forth in SEQ ID NOs: 2, 4, 6 or 8;
- 25 (b) a nucleotide sequence encoding a polypeptide as set forth in SEQ ID NOs: 10, 12 or 14 with at least one conservative amino acid substitution, wherein the polypeptide has an activity of the mature form of a polypeptide as set forth in SEQ ID NOs: 10, 12 or 14;
- 30 (c) a nucleotide sequence encoding a polypeptide as set forth in SEQ ID NOs: 2, 4, 6 or 8 with at least

one amino acid insertion, wherein the polypeptide has an activity of the mature form of a polypeptide as set forth in SEQ ID NOs: 2, 4, 6 or 8;

- (d) a nucleotide sequence encoding a polypeptide as set forth in SEQ ID NOs: 10, 12 or 14 with at least one amino acid insertion, wherein the polypeptide has an activity of the mature form of a polypeptide as set forth in SEQ ID NOs: 10, 12 or 14;
- (e) a nucleotide sequence encoding a polypeptide 10 as set forth in SEQ ID NOs: 2, 4, 6 or 8 with at least one amino acid deletion, wherein the polypeptide has an activity of the mature form of a polypeptide as set forth in SEQ ID NOs: 2, 4, 6 or 8;
- (f) a nucleotide sequence encoding a polypeptide 15 as set forth in SEQ ID NOs: 10, 12 or 14 with at least one amino acid deletion, wherein the polypeptide has an activity of the mature form of a polypeptide as set forth in SEQ ID NOs: 10, 12 or 14;
- (g) a nucleotide sequence encoding a polypeptide 20 as set forth in SEQ ID NOs: 2, 4, 6 or 8 which has a Cand/or N- terminal truncation, wherein the polypeptide has an activity of the mature form of a polypeptide as set forth in SEQ ID NOs: 2, 4, 6 or 8;
- (h) a nucleotide sequence encoding a polypeptide 25 as set forth in SEQ ID NOs: 10, 12 or 14 which has a Cand/or N- terminal truncation, wherein the polypeptide has an activity of the mature form of a polypeptide as set forth in SEQ ID NOs: 10, 12 or 14;
- (i) a nucleotide sequence encoding a polypeptide 30 as set forth in SEQ ID NOs: 2, 4, 6 or 8 with at least one modification selected from the group consisting of

10

20

25

at least one amino acid substitution, amino acid insertion, amino acid deletion, C-terminal truncation, and N-terminal truncation, wherein the polypeptide has an activity of the mature form of a polypeptide as set forth in SEQ ID NOs: 2, 4, 6 or 8;

- (j) a nucleotide sequence encoding a polypeptide as set forth in SEQ ID NOs: 10, 12 or 14 with at least one modification selected from the group consisting of at least one amino acid substitution, amino acid insertion, amino acid deletion, C-terminal truncation, and N-terminal truncation, wherein the polypeptide has an activity of the mature form of a polypeptide as set forth in SEQ ID NOs: 10, 12 or 14;
- (k) a nucleotide sequence of (a)-(j) comprising a
 15 fragment of at least about 16 nucleotides;
 - (1) a nucleotide sequence which hybridizes under moderately or highly stringent conditions to the complement of any of (a), (c), (e), (g), (i) or (k), wherein the polypeptide has an activity of the mature form of a polypeptide as set forth in SEQ ID NOs: 2, 4, 6 or 8;
 - (m) a nucleotide sequence which hybridizes under moderately or highly stringent conditions to the complement of any of (b), (d), (f), (h), (j) or (k), wherein the polypeptide has an activity of the mature form of a polypeptide as set forth in SEQ ID NOs: 10, 12 or 14; and
 - (n) a nucleotide sequence complementary to any of (a)-(m).

The invention also provides for an isolated polypeptide comprising the amino acid sequence selected from the group consisting of:

- (a) an amino acid sequence comprising the mature form of the polypeptide of SEQ ID NO: 2, and optionally further comprising an amino-terminal methionine;
 - (b) an amino acid sequence comprising the mature form of the polypeptide of SEQ ID NO: 4, and optionally further comprising an amino-terminal methionine;
- 10 (c) an amino acid sequence comprising the mature form of the polypeptide of SEQ ID NO: 6, and optionally further comprising an amino-terminal methionine;
- (d) an amino acid sequence comprising the mature form of the polypeptide of SEQ ID NO: 8, and optionally15 further comprising an amino-terminal methionine;
 - (e) an amino acid sequence comprising the mature form of the polypeptide of SEQ ID NO: 10, and optionally further comprising an amino-terminal methionine;
- 20 (f) an amino acid sequence comprising the mature form of the polypeptide of SEQ ID NO: 12, and optionally further comprising an amino-terminal methionine;
- (g) an amino acid sequence comprising the mature 25 form of the polypeptide of SEQ ID NO: 14, and optionally further comprising an amino-terminal methionine;
- (h) an amino acid sequence for an ortholog of any one of SEQ ID NOs: 2, 4, 6 or 8, wherein the encoded
 30 polypeptide has an activity of the mature form of a polypeptide as set forth in SEQ ID NOs: 2, 4, 6 or 8;

A-692

- (i) an amino acid sequence for an ortholog of any one of SEQ ID NOs: 10, 12 or 14, wherein the encoded polypeptide has an activity of the mature form of a polypeptide as set forth in SEQ ID NOs: 10, 12 or 14;
- (j) an amino acid sequence that is at least about 70, 80, 85, 90, 95, 96, 97, 98 or 99 percent identical to the amino acid sequence of SEQ ID NOs: 2, 4, 6 or 8, wherein the polypeptide has an activity of the mature form of a polypeptide as set forth in SEQ ID NOs: 2, 4, 10 6 or 8;
- (k) an amino acid sequence that is at least about 70, 80, 85, 90, 95, 96, 97, 98 or 99 percent identical to the amino acid sequence of SEQ ID NOs: 10, 12 or 14, wherein the polypeptide has an activity of the mature form of a polypeptide as set forth in SEQ ID NOs: 10, 12 or 14;
- (1) a fragment of the amino acid sequence as set forth in SEQ ID NOs: 2, 4, 6 or 8 comprising at least about 25 amino acid residues, wherein the polypeptide 20 has an activity of the mature form of a polypeptide as set forth in SEQ ID NOs: 2, 4, 6 or 8;
 - (m) a fragment of the amino acid sequence as set forth in SEQ ID NOs: 10, 12 or 14 comprising at least about 25 amino acid residues, wherein the polypeptide has an activity of the mature form of a polypeptide as set forth in SEQ ID NOs: 10, 12 or 14;
- (n) an amino acid sequence for an allelic variant or splice variant of either the amino acid sequence as set forth in SEQ ID NOs: 2, 4, 6 or 8, or at least one
 30 of (a), (c), (e), (f), (h), (i), (k) or (l), above, wherein the polypeptide has an activity of the mature form of a polypeptide as set forth in SEQ ID NOs: 2, 4,

30

A-692

6 or 8; and

(o) an amino acid sequence for an allelic variant or splice variant of either the amino acid sequence as set forth in SEQ ID NOs: 10, 12 or 14, or at least one of (b), (d), (f), (h), (j), (l) or (m) wherein the polypeptide has an activity of the mature form of a polypeptide as set forth in SEQ ID NOs: 10, 12 or 14.

The invention further provides for an isolated polypeptide comprising the amino acid sequence selected from the group consisting of:

- (a) the amino acid sequence as set forth in SEQ ID NOs: 2, 4, 6 or 8 with at least one conservative amino acid substitution, wherein the polypeptide has an activity of the mature form of a polypeptide as set forth in SEQ ID NOs: 2, 4, 6 or 8;
- (b) the amino acid sequence as set forth in SEQ ID NOs: 10, 12 or 14 with at least one conservative amino acid substitution, wherein the polypeptide has an activity of the mature form of a polypeptide as set forth in SEQ ID NOs: 10, 12 or 14;
- (c) the amino acid sequence as set forth in SEQ ID NOs: 2, 4, 6 or 8 with at least one amino acid insertion, wherein the polypeptide has an activity of the mature form of a polypeptide as set forth in SEQ ID NOs: 2, 4, 6 or 8;
 - (d) the amino acid sequence as set forth in SEQ ID NOs: 10, 12 or 14 with at least one amino acid insertion, wherein the polypeptide has an activity of the mature form of a polypeptide as set forth in SEQ ID NOs: 10, 12 or 14;

5

10

15

20

25

- (e) the amino acid sequence as set forth in SEQ ID NOs: 2, 4, 6 or 8 with at least one amino acid deletion, wherein the polypeptide has an activity of the mature form of a polypeptide as set forth in SEQ ID NOs: 2, 4, 6 or 8;
- (f) the amino acid sequence as set forth in SEQ ID NOs: 10, 12 or 14 with at least one amino acid deletion, wherein the polypeptide has an activity of the mature form of a polypeptide as set forth in SEQ ID NOs: 10, 12 or 14;
- (g) the amino acid sequence as set forth in SEQ ID NOs: 2, 4, 6 or 8 which has a C- and/or N-terminal truncation, wherein the polypeptide has an activity of the mature form of_a polypeptide as set forth in SEQ ID NOs: 2, 4, 6 or 8;
- (h) the amino acid sequence as set forth in SEQ ID NOs: 10, 12 or 14 which has a C- and/or N-terminal truncation, wherein the polypeptide has an activity of the mature form of a polypeptide as set forth in SEQ ID NOs: 10, 12 or 14;
- (i) the amino acid sequence as set forth in SEQ ID NOs: 2, 4, 6 or 8, with at least one modification selected from the group consisting of at least one amino acid substitution, amino acid insertion, amino acid deletion, C-terminal truncation, and N-terminal truncation, wherein the polypeptide has an activity of the mature form of a polypeptide as set forth in SEQ ID NOs: 2, 4, 6 or 8; and
- (j) the amino acid sequence as set forth in SEQ ID 30 NOs: 10, 12 or 14, with at least one modification selected from the group consisting of at least one amino acid substitution, amino acid insertion, amino

10

acid deletion, C-terminal truncation, and N-terminal truncation, wherein the polypeptide has an activity of the mature form of a polypeptide as set forth in SEQ ID NOs: 10, 12 or 14.

Also provided are fusion polypeptides comprising the amino acid sequences of (a)-(j) above.

The present invention also provides for an expression vector comprising the isolated nucleic acid molecules as set forth herein, recombinant host cells comprising recombinant nucleic acid molecules as set forth herein, and a method of producing a B7-like polypeptide comprising culturing the host cells and optionally isolating the polypeptide so produced.

A transgenic non-human animal comprising a nucleic acid molecule encoding a B7-like polypeptide is also encompassed by the invention. The B7-like nucleic acid molecules are introduced into the animal in a manner that allows expression and increased levels of the B7-like polypeptide, which may include increased circulating levels. The transgenic non-human animal is preferably a mammal.

Also provided are derivatives of the B7-like polypeptides of the present invention.

Additionally provided are selective binding agents such as antibodies and peptides capable of specifically binding the at least one of the B7-like polypeptides of the invention. Such antibodies and peptides may be agonistic or antagonistic.

Pharmaceutical compositions comprising the 30 nucleotides, polypeptides, or selective binding agents of the present invention and one or more

5

15

20

pharmaceutically acceptable formulation agents are also encompassed by the invention. The pharmaceutical compositions are used to provide therapeutically effective amounts of the nucleotides or polypeptides of the present invention. The invention is also directed to methods of using the polypeptides, nucleic acid molecules, and selective binding agents.

The B7-like polypeptides and nucleic acid molecules of the present invention may be used to treat, prevent, ameliorate, and/or detect diseases and disorders, including those recited herein.

The present invention also provides a method of assaying test molecules to identify a test molecule which binds to at least one B7-like polypeptide. method comprises contacting a B7-like polypeptide with a test molecule and determining the extent of binding of the test molecule to the polypeptide. The method further comprises determining whether such molecules are agonists or antagonists of at least one B7-like polypeptide. The present invention further provides a method of testing the impact of molecules on the expression of at least one B7-like polypeptide or on the activity of the mature form of at least one B7like polypeptide.

25 Methods of regulating expression and modulating (i.e., increasing or decreasing) levels of at least one B7-like polypeptide are also encompassed by One method comprises administering to an invention. animal a nucleic acid molecule encoding a B7-like 30 polypeptide. another method, a nucleic In molecule comprising elements that regulate or modulate the expression of a B7-like polypeptide may be

administered. Examples of these methods include gene therapy, cell therapy, and anti-sense therapy as further described herein.

In another aspect of the present invention, the 5 B7-like polypeptides may be used for identifying receptors thereof ("B7-like receptors"). Various forms of "expression cloning" have been extensively used for cloning receptors for protein ligands. See for example, H. Simonsen and H.F. Lodish, Trends in 10 Pharmacological Sciences, 15:437-441 (1994), and Tartaglia et al., *Cell*, <u>83</u>:1263-1271 (1995). The isolation of the B7-like receptor(s) is useful for identifying ordeveloping novel agonists and antagonists of the B7-like polypeptide-signaling 15 pathway. Such agonists and antagonists include soluble B7-like receptor(s), anti-B7-like receptor(s) selective binding agents (such as antibodies and derivatives thereof), smal1 molecules, and antisense oligonucleotides, any of which can be used for treating 20 one or more of the diseases or disorders, including those recited herein.

Brief Description of the Figures

Figure 1 depicts a nucleic acid sequence (SEQ ID NO:1) encoding human B7-like protein (B71.h1). Also depicted is the amino acid sequence (SEQ ID NO:2) of human B7-like protein (B71.h1).

Figure 2 depicts a nucleic acid sequence (SEQ ID NO:3) encoding human B7-like protein (B71.h2). Also depicted is the amino acid sequence (SEQ ID NO:4) of human B7-like protein (B71.h2).

5

15

20

25

Figure 3 depicts a nucleic acid sequence (SEQ ID NO:5) encoding human B7-like protein (B71.h3). Also depicted is the amino acid sequence (SEQ ID NO:6) of human B7-like protein (B71.h3).

Figure 4 depicts a nucleic acid sequence (SEQ ID NO:7) encoding human B7-like protein (B71.h4). Also depicted is the amino acid sequence (SEQ ID NO:8) of human B7-like protein (B71.h4).

Figure 5 depicts a nucleic acid sequence (SEQ ID NO:9) encoding murine B7-like protein (B71.m1). Also depicted is the amino acid sequence (SEQ ID NO:10) of murine B7-like protein (B71.m1).

Figure 6 depicts a nucleic acid sequence (SEQ ID NO:11) encoding murine B7-like protein B71.m2). Also depicted is the amino acid sequence (SEQ ID NO:12) of murine B7-like protein (B71.m2).

Figure 7 depicts a nucleic acid sequence (SEQ ID NO:13) encoding murine B7-like protein (B71.m3). Also depicted is the amino acid sequence (SEQ ID NO:14) of murine B7-like protein (B71.m3).

Figure 8 depicts a comparison of the amino acid sequences of SEQ ID NO: 8 $\,$ and rat B7-1 (SEQ ID NO:15).

30

Figure 9 depicts a comparison of the amino acid sequences of SEQ ID NO: 2 and SEQ ID NO:9.

30

A-692

Detailed Description of the Invention

The section headings used herein are for organizational purposes only and are not to be construed as limiting the subject matter described. All references cited in this application are expressly incorporated by reference herein.

Definitions

The terms "B7-like gene(s)" or "B7-like nucleic acid molecule(s)" or "B7-like polynucleotide(s)" refer 10 to one or more nucleic acid molecules comprising or consisting of nucleotide sequences as set forth in SEQ ID NOs: 1, 3, 5 or 7, which encode recombinant human proteins, nucleotide sequences encoding recombinant 15 human polypeptides as set forth in SEQ ID NOs: 2, 4, 6 or 8, nucleotide sequences of the DNA inserts in B71.h1, B71.h2, B71.h3 or B71.h4, and of nucleotide sequences as set forth in SEQ ID NOs: 10, 12 or 14, which encode recombinant murine proteins, nucleotide sequences encoding recombinant murine polypeptides as 20 set forth in SEQ ID NOs: 10, 12 or 14, and nucleic acid molecules as defined herein.

The term "B7-like polypeptide(s)" refers to one or more recombinant human polypeptides comprising the amino acid sequence of SEQ ID NOs: 2, 4, 6 or 8, related polypeptides thereto, or recombinant murine polypeptides comprising the amino acid sequence of SEQ ID NOs: 10, 12 or 14, and related polypeptides thereto. Related polypeptides include: B7-like polypeptide allelic variants, B7-like polypeptide orthologs, B7-like polypeptide splice variants, B7-like polypeptide variants and B7-like polypeptide derivatives. B7-like polypeptides may be mature polypeptides, as defined

25

30

35

A-692

herein, and may or may not have an amino terminal methionine residue, depending on the method by which they are prepared.

The term "B7-like polypeptide allelic variant(s)"

5 refers to one or more of several possible naturally occurring alternate forms of a gene occupying a given locus on a chromosome of an organism or a population of organisms.

The term "B7-like polypeptide derivative(s)"

10 refers to one or more of the polypeptides as set forth in SEQ ID NOs: 2, 4, 6, 8, 10, 12, or 14, B7-like polypeptide allelic variants, B7-like polypeptide orthologs, B7-like polypeptide splice variants, or B7-like polypeptide variants, as defined herein, that have been chemically modified.

The term "B7-like polypeptide fragment(s)" refers to one or more polypeptides that comprise a truncation the amino terminus (with or without a sequence) and/or a truncation at the carboxy terminus of the polypeptides as set forth in SEQ ID NOs: 2, 4, 6, 8, 10, 12 or 14, B7-like polypeptide allelic variants, B7-like polypeptide orthologs, polypeptide splice variants and/or B7-like polypeptide variant having one or more amino acid additions or substitutions or internal deletions (wherein the resulting polypeptide is at least 6 amino acids or more in length) as compared to the B7-like polypeptide amino acid sequences set forth in SEQ ID NOs: 2, 4, 6, 8, 10, 12 or 14. B7-like polypeptide fragments may result from alternative RNA splicing or vivo protease activity. In preferred embodiments, truncations comprise about 10 amino acids, or about 20 amino acids, or about 50 amino acids, or about 75 amino acids, or about 100 amino acids, or more than about 100 amino acids. The polypeptide fragments

A-692

so produced will comprise about 25 contiguous amino acids, or about 50 amino acids, or about 75 amino acids, or about 100 amino acids, or about 150 amino acids, or about 200 amino acids. Such B7-like polypeptide fragments may optionally comprise an amino terminal methionine residue. It will be appreciated that such fragments can be used, for example, to generate antibodies to B7-like polypeptides.

The term "B7-like fusion polypeptide(s)" refers to 10 or more amino acids one (such heterologous peptide or polypeptide) at the amino or carboxy terminus of one or more of the polypeptides set forth in SEQ ID NO: 2, 4, 6, 8, 10, 12 or 14, B7-like polypeptide allelic variants, B7-like polypeptide 15 orthologs, B7-like polypeptide splice variants, or B7like polypeptide variants having one or more amino acid deletions, substitutions or internal additions compared to a B7-like polypeptide amino acid sequence as set forth in SEQ ID NOs: 2, 4, 6, 8, 10, 12 or 14.

The term "B7-like polypeptide ortholog(s)" refers to one or more polypeptides from other species that correspond to the amino acid sequences as set forth in SEQ ID NOs: 2, 4, 6, 8, 10, 12 or 14. For example, mouse and human B7-like polypeptides are considered orthologs of each other.

The term "B7-like polypeptide splice variant(s)" refers to one ore more nucleic acid molecules, usually RNA, which are generated by alternative processing of intron sequences in an RNA transcript of the B7-like polypeptide amino acid sequences as set forth in SEQ ID NOs: 2, 4, 6, 8, 10, 12 or 14.

The term "B7-like polypeptide variant(s)" refers to B7-like polypeptides comprising amino acid sequences

having one or more amino acid sequence substitutions, deletions (such as internal deletions and/or B7-like polypeptide fragments), and/or additions (such internal additions and/or B7-like fusion polypeptides) 5 as compared to the B7-like polypeptide amino acid sequence set forth in SEQ ID NOs: 2, 4, 6, 8, 10, 12 or 14 (with or without a leader sequence). Variants may naturally occurring (e.g., B7-like polypeptide allelic variants, B7-like polypeptide orthologs and B7-10 polypeptide splice variants) or artificially constructed. Such B7-like polypeptide variants may be prepared from the corresponding nucleic acid molecules having a DNA sequence that varies accordingly from the DNA sequence as set forth in SEQ 15 ID NOs: 1, 3, 5, 7, 9, 11 or 13. In preferred embodiments, the variants have from 1 to 3, or from 1 to 5, or from 1 to 10, or from 1 to 15, or from 1 to 20, or from 1 to 25, or from 1 to 50, or from 1 to 75, from 1 to 100, or more than 100 amino substitutions, insertions, additions and/or deletions, 20 wherein the substitutions may be conservative, or nonconservative, or any combination thereof.

The term "antigen(s)" refers to one or more molecules or portion thereof capable of being bound by a selective binding agent, such as an antibody, and additionally capable of being used in an animal to produce antibodies capable of binding to an epitope of that antigen. An antigen may have one or more epitopes.

30 The term "biologically active B7-like polypeptide(s)" refers to one ormore B7-like polypeptides having at least one activity

A-692

characteristic of a polypeptide comprising the amino acid sequence of SEQ ID NOs: 2, 4, 6, 8, 10, 12 or 14.

The terms "effective amount" and "therapeutically effective amount" each refer to the amount of a B7-like polypeptide or B7-like nucleic acid molecule used to support an observable level of one or more biological activities of the B7-like polypeptides as set forth herein.

The term "expression vector" refers to a vector

which is suitable for use in a host cell and contains nucleic acid sequences which direct and/or control the expression of heterologous nucleic acid sequences. Expression includes, but is not limited to, processes such as transcription, translation, and RNA splicing, if introns are present.

The term "host cell" is used to refer to a cell which has been transformed, or is capable of being transformed with a nucleic acid sequence and then of expressing a selected gene of interest. The term includes the progeny of the parent cell, whether or not the progeny is identical in morphology or in genetic make-up to the original parent, so long as the selected gene is present.

The term "identity" as known in the art, refers to
25 a relationship between the sequences of two or more
polypeptide molecules or two or more nucleic acid
molecules, as determined by comparing the sequences.
In the art, "identity" also means the degree of
sequence relatedness between nucleic acid molecules or
30 polypeptides, as the case may be, as determined by the
match between strings of two or more nucleotide or two

5

10

15

20

25

30

or more amino acid sequences. "Identity" measures the percent of identical matches between the smaller of two or more sequences with gap alignments (if any) addressed by a particular mathematical model or computer program (i.e., "algorithms").

The term "similarity" is a related concept, but in contrast "identity", refers to to measure a similarity which includes both identical matches and conservative substitution matches. If two polypeptide sequences have, for example, 10/20 identical amino and the remainder are all non-conservative acids, substitutions, then the percent identity and similarity would both be 50%. If in the same example, there are 5 positions where there are conservative substitutions, then the percent identity remains 50%, but the per cent similarity would be 75% (15/20). Therefore, in cases where there are conservative substitutions, the degree of similarity between two polypeptides will be higher than the percent identity between those two polypeptides.

The term "isolated nucleic acid molecule" refers to a nucleic acid molecule of the invention that (1) has been separated from at least about 50 percent of proteins, lipids, carbohydrates or other materials with which it is naturally found when total DNA is isolated from the source cells, (2) is not linked to all or a portion of a polynucleotide to which the "isolated nucleic acid molecule" is linked in nature, operably linked to a polynucleotide which it is not linked to in nature, or (4) does not occur in nature as part of a larger polynucleotide sequence. Preferably, the isolated nucleic acid molecule of the present invention is substantially free from any other

5

10

15

20

25

30

contaminating nucleic acid molecule(s) or other contaminants that are found in its natural environment that would interfere with its use in polypeptide production or its therapeutic, diagnostic, prophylactic or research use.

The term "isolated polypeptide" refers polypeptide of the present invention that (1) has been separated from at least about percent 50 polynucleotides, lipids, carbohydrates orother materials with which it is naturally found isolated from the source cell, (2) is not linked (by covalent or noncovalent interaction) to all portion of a polypeptide to which the "isolated polypeptide" is linked in nature, (3) is operably linked (by covalent or noncovalent interaction) to a polypeptide with which it is not linked in nature, or (4) does not occur in nature. Preferably, the isolated polypeptide is substantially free from any other contaminating polypeptides or other contaminants that found in its natural are environment that would interfere with its therapeutic, diagnostic, prophylactic or research use.

The term "mature B7-like polypeptide(s)" refers to one or more B7-like polypeptide lacking a leader sequence. A mature B7-like polypeptide may also include other modifications such as proteolytic processing of the amino terminus (with or without a leader sequence) and/or the carboxy terminus, cleavage of a smaller polypeptide from a larger precursor, N-linked and/or O-linked glycosylation, and the like.

The term "nucleic acid sequence(s)" or "nucleic acid molecule(s)" refers to one or more DNA or RNA sequences. The term encompasses molecules formed from

30

A-692

any of the known base analogs of DNA and RNA such as, but not limited to 4-acetylcytosine, 8-hydroxy-N6methyladenosine, aziridinyl-cytosine, pseudoisocytosine, 5-(carboxyhydroxylmethyl) uracil, 5-5 fluorouracil, 5-bromouracil, carboxymethylaminomethyl-2-thiouracil, 5-carboxymethylaminomethyluracil, dihydrouracil, inosine, iso-pentenyladenine, 1-methyladenine, 1methylpseudouracil, 1-methylguanine, 1-methylinosine, 10 2,2-dimethyl-guanine, 2-methyladenine, 2-methylguanine, 3-methylcytosine, 5-methylcytosine, N6-methyladenine, 7-methylguanine, 5-methylaminomethyluracil, 5methoxyamino-methyl-2-thiouracil, beta-Dmannosylqueosine, 5' -methoxycarbonyl-methyluracil, 5-15 methoxyuracil, 2-methylthio-N6-isopentenyladenine, uracil-5-oxyacetic acid methylester, uracil-5-oxyacetic oxybutoxosine, pseudouracil, acid, queosine, thiocytosine, 5-methyl-2-thiouracil, 2-thiouracil, 4thiouracil, 5-methyluracil, N-uracil-5-oxyacetic acid 20 methylester, uracil-5-oxyacetic acid, pseudouracil, queosine, 2-thiocytosine, and 2,6-diaminopurine.

The term "naturally occurring" or "native" when used in connection with biological materials such as nucleic acid molecules, polypeptides, host cells, and the like, refers to materials which are found in nature and are not manipulated by man. Similarly, "non-naturally occurring" or "non-native" as used herein refers to a material that is not found in nature or that has been structurally modified or synthesized by man.

The term "operably linked" is used herein to refer to an arrangement of flanking sequences wherein the

A-692

flanking sequences so described are configured or assembled so as to perform their usual function. flanking sequence operably linked to a coding sequence may be capable of effecting the replication, 5 transcription and/or translation of the For example, a coding sequence is operably sequence. linked to a promoter when the promoter is capable of directing transcription of that coding sequence. flanking sequence need not be contiguous with the 10 coding sequence, so long as it functions correctly. Thus, for example, intervening untranslated transcribed sequences can be present between a promoter sequence and the coding sequence and the promoter sequence can still be considered "operably linked" to 15 the coding sequence.

The term "pharmaceutically acceptable carrier(s)" or "physiologically acceptable carrier(s)" as used herein refers to one or more formulation materials suitable for accomplishing or enhancing the delivery of the B7-like polypeptide, B7-like nucleic acid molecule or B7-like selective binding agent as a pharmaceutical composition.

The term "selective binding agent(s)" refers to a molecule or molecules having specificity for a B7-like polypeptide. As used herein, the terms, "specific" and 25 "specificity" refer to the ability of the selective binding agents to bind to human B7-like polypeptides and not to bind to human non-B7-like polypeptides. will be appreciated, however, that the selective 30 binding agents may also bind orthologs οf polypeptides as set forth in SEQ ID NOs: 2, 4, 6, 8, 10, 12 or 14, that is, interspecies versions thereof,

5

such as mouse and rat polypeptides.

The term "transduction" is used to refer to the transfer of genes from one bacterium to another, usually by a phage. "Transduction" also refers to the acquisition and transfer of eukaryotic cellular sequences by retroviruses.

The term "transfection" is used to refer to the uptake of foreign or exogenous DNA by a cell, and a cell has been "transfected" when the exogenous DNA has been introduced inside the cell membrane. A number of 10 transfection techniques are well known in the art and are disclosed herein. See, for example, Graham et al., Virology, 52:456 (1973); Sambrook et al., Molecular Cloning, a laboratory Manual, Cold Spring Harbor Laboratories New York (1989); Davis et al., 15 Methods in Molecular Biology, Elsevier (1986); and Chu et al., Gene, 13:197 (1981). Such techniques can be used to introduce one or more exogenous DNA moieties into suitable host cells.

The term "transformation" as used herein refers to a change in a cell's genetic characteristics, and a cell has been transformed when it has been modified to contain a new DNA. For example, a cell is transformed where it is genetically modified from its native state.

Following transfection or transduction, the transforming DNA may recombine with that of the cell by physically integrating into a chromosome of the cell, may be maintained transiently as an episomal element without being replicated, or may replicate

30 independently as a plasmid. A cell is considered to

5

10

15

have been stably transformed when the DNA is replicated with the division of the cell.

The term "vector" is used to refer to any molecule (e.g., nucleic acid, plasmid, or virus) used to transfer coding information to a host cell.

Relatedness of Nucleic Acid Molecules and/or Polypeptides

It is understood that related nucleic acid molecules include allelic or splice variants of the nucleic acid molecules of SEQ ID NOs: 1, 3, 5, 7, 9, 11 or 13, and include sequences which are complementary to any of the above nucleotide sequences. Related nucleic acid molecules also include a nucleotide sequence encoding a polypeptide comprising or consisting essentially of a substitution, modification, addition and/or a deletion of one or more amino acid residues compared to the polypeptides in SEQ ID NOs: 2, 4, 6, 8, 10, 12 or 14.

Fragments include molecules which encode a 20 polypeptide of at least about 25 amino acid residues, or about 50, or about 75, or about 100, or greater than about 100 amino acid residues of the polypeptides of SEQ ID NOs: 2, 4, 6, 8, 10, 12 or 14.

addition, In related B7-like nucleic acid 25 molecules include those molecules which comprise nucleotide sequences which hybridize under moderately or highly stringent conditions as defined herein with the fully complementary sequence of the nucleic acid molecules of SEQ ID NOs: 1, 3, 5, 7, 9, 11 or 13, or of 30 a molecule encoding a polypeptide, which polypeptide

5

10

15

20

25

comprises the amino acid sequence as shown in SEQ ID NOs: 2, 4, 6, 8, 10, 12 or 14, or of a nucleic acid fragment as defined herein, or of a nucleic acid fragment encoding a polypeptide as defined herein. Hybridization probes may be prepared using the B7-like sequences provided herein to screen cDNA, genomic or synthetic DNA libraries for related sequences. Regions of the DNA and/or amino acid sequence of B7-like polypeptides that exhibit significant identity to known sequences are readily determined using sequence alignment algorithms as described herein, and those regions may be used to design probes for screening.

The term "highly stringent conditions" refers to those conditions that are designed to permit hybridization of DNA strands whose sequences are highly complementary, and to exclude hybridization significantly mismatched Hybridization DNAs. stringency is principally determined by temperature, ionic strength, and the concentration of denaturing agents such as formamide. Examples of "highly stringent conditions" for hybridization and washing are 0.015M sodium chloride, 0.0015M sodium citrate at 65-68°C or 0.015M sodium chloride, 0.0015M sodium citrate, and 50% formamide at 42°C. See Sambrook, Fritsch & Maniatis, Molecular Cloning: A Laboratory Manual, Ed., Cold Spring Harbor Laboratory, (Cold Harbor, N.Y. 1989); Anderson et al., Nucleic Acid Hybridisation: a practical approach, Ch. 4, IRL Press Limited (Oxford, England).

More stringent conditions (such as higher temperature, lower ionic strength, higher formamide, or other denaturing agent) may also be used, however, the

rate of hybridization will be affected. Other agents may be included in the hybridization and washing buffers for the purpose of reducing non-specific and/or background hybridization. Examples are 0.1% bovine 5 serum albumin, 0.1% polyvinyl-pyrrolidone, 0.1% sodium pyrophosphate, 0.1% sodium dodecylsulfate (NaDodSO4 or ficoll, Denhardt's solution, sonicated salmon SDS), sperm DNA (or other non-complementary DNA), and dextran sulfate, although other suitable agents can also be 10 used. The concentration and types of these additives can be changed without substantially affecting the stringency of the hybridization conditions. Hybridization experiments are usually carried out at pH 6.8-7.4, however, at typical ionic strength conditions, 15 the rate of hybridization is nearly independent of pH. See Anderson et al., Nucleic Acid Hybridisation: a Practical Approach, Ch. 4, IRL Press Limited (Oxford, England).

Factors affecting the stability of a DNA duplex 20 include base composition, length, and degree of base pair mismatch. Hybridization conditions can be adjusted by one skilled in the art in order to accommodate these variables and allow DNAs of different sequence relatedness to form hybrids. The melting 25 temperature of a perfectly matched DNA duplex can be estimated by the following equation:

$$T_m(^{\circ}C) = 81.5 + 16.6(log[Na+]) + 0.41(%G+C) - 600/N - 0.72(%formamide)$$

where N is the length of the duplex formed, [Na+] 30 is the molar concentration of the sodium ion in the hybridization or washing solution, %G+C is the percentage of (guanine+cytosine) bases in the hybrid.

For imperfectly matched hybrids, the melting temperature is reduced by approximately 1°C for each 1% mismatch.

The term "moderately stringent conditions" refers to conditions under which a DNA duplex with a greater 5 degree of base pair mismatching than could occur under "highly stringent conditions" is able to form. Examples of typical "moderately stringent conditions" are 0.015M sodium chloride, 0.0015M sodium citrate at 10 50-65°C or 0.015M sodium chloride, 0.0015M sodium citrate, and 20% formamide at 37-50°C. By way of example, a "moderately stringent" condition of 50°C in 0.015 M sodium ion will allow about a 21% mismatch.

It will be appreciated by those skilled in the art 15 that there is no absolute distinction between "highly" and "moderately" stringent conditions. For example, at 0.015M sodium ion formamide), (no the melting temperature of perfectly matched long DNA is about 71°C. With a wash at 65° C (at the same 20 strength), this would allow for approximately a 6% mismatch. To capture more distantly related sequences, one skilled in the art can simply lower the temperature or raise the ionic strength.

A good estimate of the melting temperature in 1M 25 NaCl* for oligonucleotide probes up to about 20nt is given by:

 $Tm = 2^{\circ}C$ per A-T base pair + $4^{\circ}C$ per G-C base pair

*The sodium ion concentration in 6X salt sodium citrate (SSC) is 1M. See Suggs et al., Developmental

5

10

15

20

Biology Using Purified Genes, p. 683, Brown and Fox (eds.) (1981).

High stringency washing conditions for oligonucleotides are usually at a temperature of $0-5^{\circ}$ C below the Tm of the oligonucleotide in 6X SSC, 0.1% SDS.

another embodiment, related nucleic In acid molecules comprise or consist of a nucleotide sequence that is about 70 percent identical to the nucleotide sequences as shown in SEQ ID NOs: 1, 3, 5, 7, 9, 11 or 13, or comprise or consist essentially of a nucleotide sequence encoding a polypeptide that is about percent identical to the polypeptides as set forth in SEQ ID NOs: 2, 4, 6, 8, 10, 12 or 14. In preferred embodiments, the nucleotide sequences are about percent, or about 80 percent, or about 85 percent, or about 90 percent, or about 95, 96, 97, 98, or percent identical to the nucleotide sequences as shown in SEQ ID NOs: 1, 3, 5, 7, 9, 11 or 13, or the nucleotide sequences encode a polypeptide that is about 75 percent, or about 80 percent, or about 85 percent, or about 90 percent, or about 95, 96, 97, 98, or 99 percent identical to the polypeptide sequences as set forth in SEQ ID NOs: 2, 4, 6, 8, 10, 12 or 14.

Differences in the nucleic acid sequence may result in conservative and/or non-conservative modifications of an amino acid sequence relative to an amino acid sequence of SEQ ID NOs: 2, 4, 6, 8, 10, 12 or 14.

30 Conservative modifications to an amino acid sequence of SEQ ID NOs: 2, 4, 6, 8, 10, 12 or 14 (and

10

the corresponding modifications to the encoding nucleotides) will produce B7-like polypeptides having functional and chemical characteristics those of naturally occurring B7-like polypeptides. contrast, substantial modifications in the functional and/or chemical characteristics of B7-like polypeptides may be accomplished by selecting substitutions in an amino acid sequence of SEQ ID NOs: 2, 4, 6, 8, 10, 12 or 14 that differ significantly in their effect on maintaining (a) the structure of the molecular backbone in the area of the substitution, for example, as a sheet or helical conformation, (b) the charge or hydrophobicity of the molecule at the target site, or (c) the bulk of the side chain.

15 For example, а "conservative amino acid substitution" may involve a substitution of a native amino acid residue with a nonnative residue such that there is little or no effect on the polarity or charge the amino acid residue at that position. 20 Furthermore, any native residue in the polypeptide may also be substituted with alanine, as has previously described for "alanine scanning mutagenesis."

Desired amino acid substitutions (whether conservative or non-conservative) can be determined by those skilled in the art at the time such substitutions are desired. For example, amino acid substitutions can be used to identify important residues of the B7-like polypeptides, or to increase or decrease the affinity of the B7-like polypeptides described herein.

Exemplary amino acid substitutions are set forth in Table I.

Table I

Amino Acid Substitutions

Original Residues	Exemplary Substitutions	Preferred Substitutions
Ala	Val, Leu, Ile	Val
Arg	Lys, Gln, Asn	Lys
Asn	Gln	Gln
Asp	Glu	Glu
Cys	Ser, Ala	Ser
Gln	Asn	Asn
Glu	Asp	Asp
Gly	Pro, Ala	Ala
His	Asn, Gln, Lys, Arg	Arg
Ile	Leu, Val, Met, Ala,	Leu
	Phe, Norleucine	
Leu	Norleucine, Ile,	Ile
	Val, Met, Ala, Phe	
Lys	Arg, 1,4 Diamino-butyric	Arg
	Acid, Gln, Asn	
Met	Leu, Phe, Ile	Leu
Phe	Leu, Val, Ile, Ala,	Leu
	Tyr	
Pro	Ala	Gly
Ser	Thr, Ala, Cys	Thr
Thr	Ser	Ser
Trp	Tyr, Phe	Tyr
Tyr	Trp, Phe, Thr, Ser	Phe
Val	Ile, Met, Leu, Phe,	Leu
	Ala, Norleucine	

⁵ Conservative amino acid substitutions also encompass non-naturally occurring amino acid residues

A-692

which are typically incorporated by chemical peptide synthesis rather than by synthesis in biological systems. These include peptidomimetics, and other reversed or inverted forms of amino acid moieties.

Naturally occurring residues may be divided into classes based on common side chain properties:

- 1) hydrophobic: norleucine, Met, Ala, Val, Leu, Ile;
- 2) neutral hydrophilic: Cys, Ser, Thr, Asn,
 Gln;
- 3) acidic: Asp, Glu;
- 4) basic: His, Lys, Arg;
- 5) residues that influence chain orientation: Gly, Pro; and
- 15 6) aromatic: Trp, Tyr, Phe.

For example, non-conservative substitutions may involve the exchange of a member of one of these classes for a member from another class. Such substituted residues may be introduced into regions of the human B7-like polypeptides that are homologous with non-human B7-like polypeptide orthologs, or into the non-homologous regions of the molecule.

In making such changes, the hydropathic index of amino acids may be considered. Each amino acid has been assigned a hydropathic index on the basis of their hydrophobicity and charge characteristics, these are: isoleucine (+4.5); valine (+4.2); leucine (+3.8); phenylalanine (+2.8); cysteine/cystine (+2.5); methionine (+1.9); alanine (+1.8); glycine (-0.4); threonine (-0.7); serine (-0.8); tryptophan (-0.9);

25

A-692

tyrosine (-1.3); proline (-1.6); histidine (-3.2); glutamate (-3.5); glutamine (-3.5); aspartate (-3.5); asparagine (-3.5); lysine (-3.9); and arginine (-4.5).

The importance of the hydropathic amino acid index 5 in conferring interactive biological function on a protein is understood in the art. Kyte et al., J. Mol. Biol., 157:105-131 (1982). It is known that certain amino acids may be substituted for other amino acids having a similar hydropathic index or score and still 10 similar biological activity. retain a In making changes based upon the hydropathic index, substitution of amino acids whose hydropathic indices are within ±2 is preferred, those which are within ±1 are particularly preferred, and those within ±0.5 are 15 even more particularly preferred.

It. is also understood in the art that the substitution of like amino acids can be made effectively on the basis of hydrophilicity, particularly where the biologically functionally equivalent protein or peptide thereby created intended for use in immunological embodiments, as in present case. The greatest local average hydrophilicity of a protein, as governed by the hydrophilicity of its adjacent amino acids, correlates with its immunogenicity and antigenicity, i.e., with a biological property of the protein.

The following hydrophilicity values have been assigned to amino acid residues: arginine (+3.0); lysine (+3.0); aspartate (+3.0 ± 1); glutamate (+3.0 ± 1); serine (+0.3); asparagine (+0.2); glutamine (+0.2); glycine (0); threonine (-0.4); proline (-0.5 ± 1);

10

alanine (-0.5); histidine (-0.5); cysteine (-1.0); methionine (-1.3); valine (-1.5); leucine (-1.8);isoleucine (-1.8); tyrosine (-2.3); phenylalanine (-2.5); tryptophan (-3.4). In making changes based upon 5 similar hydrophilicity values, the substitution of amino acids whose hydrophilicity values are within ±2 preferred, those which are within particularly preferred, and those within ±0.5 are even more particularly preferred. One may also identify epitopes from primary amino acid sequences on the basis of hydrophilicity. These regions are also referred to as "epitopic core regions."

A skilled artisan will be able to determine suitable variants of the polypeptide as set forth in 15 SEQ ID NOs: 2, 4, 6, 8, 10, 12 or 14 using well-known techniques. For identifying suitable areas of the molecule that may be changed without destroying activity, one skilled in the art may target areas not believed to be important for activity. For example, 20 when similar polypeptides with similar activities from the same species or from other species are known, one skilled in the art may compare the amino acid sequence of a B7-like polypeptide to such similar polypeptides. 25 With such a comparison, one can identify residues and portions of the molecules that are conserved among similar polypeptides. It will be appreciated that changes in areas of a B7-like polypeptide that are not conserved relative to such similar polypeptides would 30 be less likely to adversely affect the biological activity and/or structure of such B7-like polypeptide. One skilled in the art would also know that, even in

5

25

30

relatively conserved regions, one may substitute chemically similar amino acids for the naturally occurring residues while retaining activity (conservative amino acid residue substitutions). Therefore, even areas that may be important for biological activity or for structure may be subject to conservative amino acid substitutions destroying the biological activity or without adversely affecting the polypeptide structure.

Additionally, one skilled in the art can review structure-function studies identifying residues in similar polypeptides that are important for activity or structure. In view of such a comparison, one can predict the importance of amino acid residues in the B7-like polypeptides that correspond to amino acid residues that are important for activity or structure in similar polypeptides. One skilled in the art may opt for chemically similar amino acid substitutions for such predicted important amino acid residues of B7-like polypeptides.

One skilled in the art can also analyze the three-dimensional structure and amino acid sequence in relation to that structure in similar polypeptides. In view of that information, one skilled in the art may predict the alignment of amino acid residues of a B7-like polypeptide with respect to its three-dimensional structure. One skilled in the art may choose not to make radical changes to amino acid residues predicted to be on the surface of the protein, since such residues may be involved in important interactions with other molecules. Moreover, one skilled in the art may generate test variants containing a single amino acid

5

10

substitution at each desired amino acid residue. variants can then be screened using activity assays know to those skilled in the art. Such variants could be used to gather information about suitable variants. For example, if one discovered that a change to a particular amino acid residue resulted in destroyed, undesirably reduced, or unsuitable activity, variants with such a change would be avoided. In other words, based on information gathered from such routine experiments, one skilled in the art can readily determine the amino acids where further substitutions should be avoided either alone or in combination with other mutations.

A number of scientific publications have been devoted to the prediction of secondary structure. 15 Moult J., Curr. Op. in Biotech., 7(4):422-427 (1996), Chou et al., Biochemistry, 13(2):222-245 (1974); Chou et al., Biochemistry, 113(2):211-222 (1974); Chou et al., Adv. Enzymol. Relat. Areas Mol. Biol., 47:45-148 (1978); Chou et al., Ann. Rev. Biochem., 47:251-276 and 20 Chou et al., Biophys. J., 26:367-384 (1979). Moreover, computer programs are currently available to assist with predicting secondary structure. One method of predicting secondary structure is based upon homology 25 modeling. For example, two polypeptides or proteins which have a sequence identity of greater than 30%, or similarity greater than 40% often have similar structural topologies. The recent growth of the protein structural data base (PDB) has provided 30 enhanced predictability of secondary structure, including the potential number of folds within a polypeptide's or protein's structure. See Holm et al.,

5

25

30

Nucl. Acid. Res., 27(1):244-247 (1999). It has been suggested (Brenner et al., Curr. Op. Struct. Biol., 7(3):369-376 (1997)) that there are a limited number of folds in a given polypeptide or protein and that once a critical number of structures have been resolved, structural prediction will gain dramatically accuracy.

Additional methods of predicting secondary structure include "threading" (Jones, D., Curr. Opin. 10 Struct. Biol., 7(3):377-387 (1997); Sippl et al., Structure, 4(1):15-9 (1996)), "profile analysis" (Bowie et al., Science, 253:164-170 (1991); Gribskov et al., Meth. Enzym., 183:146-159 (1990); Gribskov et al., Proc. Nat. Acad. Sci., 84(13):4355-4358 (1987)), and "evolutionary linkage" (See Holm, supra, and Brenner, 15 supra).

Preferred B7-like polypeptide variants include glycosylation variants wherein the number and/or type of glycosylation sites have been altered compared to a 20 amino acid sequence set forth in SEQ ID NOs: 2, 4, 6, 8, 10, 12 or 14. In one embodiment, polypeptide variants comprise a greater or a lesser number of N-linked glycosylation sites than an amino acid sequence set forth in SEQ ID NOs: 2, 4, 6, 8, 10, 12 14. N-linked glycosylation An site characterized by the sequence Asn-X-Ser or Asn-X-Thr, wherein the amino acid residue designated as X may be any amino acid residue except proline. substitution(s) of amino acid residues to create this sequence provides a potential new site for the addition of an N-linked carbohydrate chain. Alternatively, substitutions which eliminate this sequence will remove

5

10

15

20

25

30

an existing N-linked carbohydrate chain. Also provided is a rearrangement of N-linked carbohydrate chains wherein one more N-linked glycosylation sites or (typically those that are naturally occurring) are eliminated and one or more new N-linked sites are created. Additional preferred B7-like variants include cysteine variants, wherein one or more cysteine residues are deleted from or substituted for another amino acid (e.g., serine) as compared to an amino acid sequences set forth in SEQ ID NOs: 2, 4, 6, 8, 10, 12 or 14. Cysteine variants are useful when B7-like polypeptides must be refolded into a biologically active conformation such as after the isolation of insoluble inclusion bodies. Cysteine variants generally have fewer cysteine residues than the native protein, and typically have an even number to minimize interactions resulting from unpaired cysteines.

Figure 8 depicts a topological comparison of the amino acid sequences of SEQ ID NO:10 and rat B7-1 (SEQ ID NO:15). Figure 9 depicts a topological comparison of the amino acid sequences of SEQ ID NO: 2 and SEQ ID NO:10. A skilled artisan can readily discern the residues conserved among the family members and, with this information, can readily generate variants within the scope of this invention, using standard techniques.

In addition, a polypeptide comprising the amino acid sequence of SEQ ID NOs: 2, 4, 6, 8, 10, 12, or 14 or a B7-like polypeptide variant may be fused to a homologous polypeptide to form a homodimer or to a heterologous polypeptide to form a heterodimer. Heterologous peptides and polypeptides include, but are not limited to: an epitope to allow for the detection

20

25

30

A-692

and/or isolation of a B7-like fusion polypeptide; a transmembrane receptor protein or a portion thereof, such as an extracellular domain, or a transmembrane and intracellular domain; a ligand or a portion thereof 5 which binds to a transmembrane receptor protein; portion thereof enzyme or which is catalytically active; а polypeptide or peptide which promotes oligomerization, such as a leucine zipper domain; a polypeptide or peptide which increases stability, such 10 as an immunoglobulin constant region; and a polypeptide which has a therapeutic activity different from the polypeptide comprising an amino acid sequence as set forth in SEQ ID NOs: 2, 4, 6, 8, 10, 12 or 14 or a B7like polypeptide variant. Specific fusions may include the fusion of one or more of SEQ ID Nos: 2, 4, 6, 8, 15 10, 12 or 14.

Fusions can be made either at the amino terminus carboxy terminus the of the polypeptide comprising an amino acid sequence set forth in SEQ ID NOs: 2, 4, 6, 8, 10, 12 or 14 or a B7-like polypeptide variant. Fusions may be direct with no linker or adapter molecule or indirect using a linker or adapter molecule. A linker or adapter molecule may be one or more amino acid residues, typically up to about 20 to about 50 amino acid residues. A linker or adapter molecule may also be designed with a cleavage site for a DNA restriction endonuclease or for a protease to allow for the separation of the fused moieties. will be appreciated that once constructed, the fusion polypeptides can be derivatized according to methods described herein.

10

15

20

In a further embodiment of the invention, polypeptide comprising an amino acid sequence of SEQ ID NOs: 2, 4, 6, 8, 10, 12 or 14 or a B7-like polypeptide variant is fused to one or more domains of an Fc region of human IgG. Antibodies comprise two functionally independent parts, a variable domain known as "Fab", which binds antigen, and a constant domain known as "Fc", which is involved in effector functions such as complement activation and attack by phagocytic cells. An Fc has a long serum half-life, whereas an Fab is short-lived. Capon et al., Nature, 337:525-531 (1989). When constructed together with a therapeutic protein, an domain can provide longer half-life incorporate such functions as Fc receptor binding, protein A binding, complement fixation and perhaps even placental transfer. Id. Table II summarizes the use of certain Fc fusions known in the art.

Table II
Fc Fusion with Therapeutic Proteins

Form of Fc	Fusion partner	Therapeutic implications	Reference
IgG1	N-terminus of CD30-L	Hodgkin's disease; anaplastic lymphoma; T- cell leukemia	U.S. Patent No. 5,480,981
Murine Fcγ2a	IL-10	anti- inflammatory; transplant rejection	Zheng et al. (1995), J. Immunol., 154: 5590-5600
IgG1	TNF receptor	septic shock	Fisher et al. (1996), N. Engl. J. Med., 334: 1697- 1702; Van Zee et al., (1996), J. Immunol., 156: 2221-2230
IgG, IgA,	TNF	inflammation,	U.S. Pat. No.

A-692

IgM, or IgE (excluding the first domain)	receptor	autoimmune disorders	5,808,029, issued September 15, 1998
IgG1	CD4 receptor	AIDS	Capon et al. (1989), Nature 337: 525-531
IgG1, IgG3	N-terminus of IL-2	anti-cancer, antiviral	Harvill et al. (1995), Immunotech., 1: 95- 105
IgG1	C-terminus of OPG	osteoarthritis; bone density	WO 97/23614, published July 3, 1997
IgG1	N-terminus of leptin	anti-obesity	PCT/US 97/23183, filed December 11, 1997
Human Ig Cγ1	CTLA-4	autoimmune disorders	Linsley (1991), <i>J. Exp. Med.</i> , 174:561-569

In one example, all or a portion of the human IgG hinge, CH2 and CH3 regions may be fused at either the N-terminus or C-terminus of the B7-like polypeptides using methods known to the skilled artisan. resulting B7-like fusion polypeptide may be purified by use of a Protein A affinity column. Peptides and proteins fused to an Fc region have been found to exhibit a substantially greater half-life in vivo than the unfused counterpart. Also, a fusion to an Fc 10 region allows for dimerization/multimerization of the fusion polypeptide. The Fc region may be a naturally occurring Fc region, or may be altered to improve certain qualities, such as therapeutic qualities, circulation time, reduce aggregation, etc.

Identity and similarity of related nucleic acid molecules and polypeptides can be readily calculated by known methods. Such methods include, but are not limited to, those described in Computational Molecular

5

10

15

20

25

Biology, Lesk, A.M., ed., Oxford University Press, New York, 1988; Biocomputing: Informatics and Genome Projects, Smith, D.W., ed., Academic Press, New York, 1993; Computer Analysis of Sequence Data, Part 1, Griffin, A.M., and Griffin, H.G., eds., Humana Press, New Jersey, 1994; Sequence Analysis in Molecular Biology, von Heinje, G., Academic Press, 1987; Sequence Analysis Primer, Gribskov, M. and Devereux, J., eds., M. Stockton Press, New York, 1991; and Carillo et al., SIAM J. Applied Math., 48:1073 (1988).

Preferred methods to determine identity and/or similarity are designed to give the largest match between the sequences tested. Methods to determine identity and similarity are described in publicly available computer programs. Preferred computer program methods to determine identity and similarity between two sequences include, but are not limited to, the GCG program package, including GAP (Devereux et al., Nucl. Acid. Res., $\underline{12}$:387 (1984); Genetics Computer Group, University of Wisconsin, Madison, WI), BLASTP, BLASTN, and FASTA (Altschul et al., J. Mol. Biol., 215:403-410 (1990)). The BLASTX program is publicly available from the National Center for Biotechnology Information (NCBI) and other sources (BLAST Manual, Altschul et al. NCB/NLM/NIH Bethesda, MD Altschul et al., supra). The well-known Smith Waterman algorithm may also be used to determine identity.

Certain alignment schemes for aligning two amino acid sequences may result in the matching of only a short region of the two sequences, and this small aligned region may have very high sequence identity even though there is no significant relationship

5

10

15

20

25

between the two full-length sequences. Accordingly, in a preferred embodiment, the selected alignment method (GAP program) will result in an alignment that spans at least 50 contiguous amino acids of the target polypeptide.

For example, using the computer algorithm GAP (Genetics Computer Group, University of Wisconsin, Madison, WI), two polypeptides for which the percent sequence identity is to be determined are aligned for optimal matching of their respective amino acids (the "matched span", as determined by the algorithm). A gap opening penalty (which is calculated as 3X the average diagonal; the "average diagonal" is the average of the diagonal of the comparison matrix being used; the "diagonal" is the score or number assigned to each perfect amino acid match by the particular comparison matrix) and a gap extension penalty (which is usually 1/10 times the gap opening penalty), as well as a comparison matrix such as PAM 250 or BLOSUM 62 are used conjunction with the algorithm. Α standard comparison matrix (see Dayhoff et al., Atlas of Protein Sequence and Structure, vol. 5, supp.3 (1978) for the PAM 250 comparison matrix; Henikoff et al., Proc. Natl. Acad. Sci USA, 89:10915-10919 (1992) for the BLOSUM 62 comparison matrix) is also used by the algorithm.

Preferred parameters for a polypeptide sequence comparison include the following:

Algorithm: Needleman et al., J. Mol. Biol., 48:443-453 (1970);

Comparison matrix: BLOSUM 62 from Henikoff et al., Proc. Natl. Acad. Sci. USA, 89:10915-10919 (1992);

Gap Penalty: 12

5 Gap Length Penalty: 4

Threshold of Similarity: 0

The GAP program is useful with the above parameters. The aforementioned parameters are the default parameters for polypeptide comparisons (along with no penalty for end gaps) using the GAP algorithm.

Preferred parameters for nucleic acid molecule sequence comparisons include the following:

Algorithm: Needleman *et al.*, *J. Mol Biol.*, <u>48</u>:443-15 453 (1970);

Comparison matrix: matches = +10, mismatch = 0

Gap Penalty: 50

Gap Length Penalty: 3

20 The GAP program is also useful with the The aforementioned parameters are the parameters. default parameters for nucleic acid molecule comparisons.

Other exemplary algorithms, gap opening penalties,

gap extension penalties, comparison matrices,
thresholds of similarity, etc. may be used, including
those set forth in the Program Manual, Wisconsin
Package, Version 9, September, 1997. The particular
choices to be made will be apparent to those of skill
in the art and will depend on the specific comparison
to be made, such as DNA to DNA, protein to protein,

protein to DNA; and additionally, whether the comparison is between given pairs of sequences (in which case GAP or BestFit are generally preferred) or between one sequence and a large database of sequences (in which case FASTA or BLASTA are preferred).

Synthesis

It will be appreciated by those skilled in the art that the nucleic acid and polypeptide molecules described herein may be produced by recombinant and other means.

Nucleic Acid Molecules

The nucleic acid molecules encoding a polypeptide comprising the amino acid sequence of a B7-like polypeptide can readily be obtained in a variety of ways including, without limitation, chemical synthesis, cDNA or genomic library screening, expression library screening and/or PCR amplification of cDNA.

Recombinant DNA methods used herein are generally those set forth in Sambrook et al., Molecular Cloning:

A Laboratory Manual, Cold Spring Harbor Laboratory Press, Cold Spring Harbor, NY (1989), and/or Ausubel et al., eds., Current Protocols in Molecular Biology,

Green Publishers Inc. and Wiley and Sons, NY (1994). The present invention provides for nucleic acid molecules as described herein and methods for obtaining the molecules.

Where a gene encoding the amino acid sequence of a 30 B7-like polypeptide has been identified from one species, all or a portion of that gene may be used as a

10

15

20

25

30

probe to identify orthologs or related genes from the same species. The probes or primers may be used to screen cDNA libraries from various tissue sources believed to express a B7-like polypeptide. In addition, part or all of a nucleic acid molecule having the sequences as set forth in SEQ ID NOs: 1, 3, 5, 7, 9, 11 or 13 may be used to screen a genomic library to identify and isolate a gene encoding the amino acid B7-like polypeptide. sequence of a Typically, conditions of moderate or high stringency will be employed for screening to minimize the number of false positives obtained from the screen.

Nucleic acid molecules encoding the amino acid sequence of B7-like polypeptides may also be identified by expression cloning which employs the detection of positive clones based upon a property of the expressed protein. Typically, nucleic acid libraries are screened by the binding of an antibody or other binding partner (e.g., receptor or ligand) to cloned proteins which are expressed and displayed on a host cell surface. The antibody or binding partner is modified with a detectable label to identify those cells expressing the desired clone.

Recombinant expression techniques conducted in accordance with the descriptions set forth below may be followed to produce these polynucleotides and to express the encoded polypeptides. For example, by inserting a nucleic acid sequence which encodes the amino acid sequence of a B7-like polypeptide into an appropriate vector, one skilled in the art can readily produce large quantities of the desired nucleotide sequence. The sequences can then be used to generate

5

10

15

20

25

30

detection probes or amplification primers. Alternatively, a polynucleotide encoding the amino acid sequence of a B7-like polypeptide can be inserted into an expression vector. By introducing the expression vector into an appropriate host, an encoded B7-like polypeptide may be produced in large amounts.

Another method for obtaining a suitable nucleic acid sequence is the polymerase chain reaction (PCR). In this method, cDNA is prepared from poly(A)+RNA or total RNA using the enzyme reverse transcriptase. Two primers, typically complementary to two separate regions of cDNA (oligonucleotides) encoding the amino acid sequence of a B7-like polypeptide, are then added to the cDNA along with a polymerase such as Taq polymerase, and the polymerase amplifies the cDNA region between the two primers.

Another means of preparing a nucleic acid molecule encoding the amino acid sequence οf а B7-like polypeptide is chemical synthesis using methods well known to the skilled artisan such as those described by Engels et al., Angew. Chem. Intl. Ed., 28:716-734 (1989).These methods include, inter alia, phosphotriester, phosphoramidite, and H-phosphonate methods for nucleic acid synthesis. A preferred method for such chemical synthesis is polymer-supported synthesis using standard phosphoramidite chemistry. Typically, the DNA encoding the amino acid sequence of B7-like polypeptide will be several nucleotides in length. Nucleic acids larger than about 100 nucleotides can be synthesized as several fragments using these methods. The fragments can then be ligated together to form the full-length nucleotide sequence of

a B7-like polypeptide. Usually, the DNA fragment encoding the amino terminus of the polypeptide will have an ATG, which encodes a methionine residue. This methionine may or may not be present on the mature form of a B7-like polypeptide, depending on whether the polypeptide produced in the host cell is designed to be secreted from that cell. Other methods known to the skilled artisan may be used as well.

certain embodiments, nucleic acid variants contain codons which have been altered for the optimal 10 expression of a B7-like polypeptide in a given host Particular codon alterations will depend upon the B7-like polypeptide(s) and host cell(s) selected for expression. Such "codon optimization" 15 carried out by a variety of methods, for example, by selecting codons which are preferred for use in highly expressed genes in a given host cell. Computer algorithms which incorporate codon frequency tables such as "Ecohigh.cod" for codon preference of highly expressed bacterial genes may be used and are provided 20 by the University of Wisconsin Package Version 9.0, Genetics Computer Group, Madison, WI. Other useful codon frequency tables include "Celegans_high.cod", "Celegans_low.cod", "Drosophila_high.cod", 25 "Human_high.cod", "Maize_high.cod", and

Vectors and Host Cells

"Yeast_high.cod".

A nucleic acid molecule encoding the amino acid sequence of a B7-like polypeptide may be inserted into an appropriate expression vector using standard ligation techniques. The vector is typically selected to be functional in the particular host cell employed

(i.e., the vector is compatible with the host cell
machinery such that amplification of the gene and/or
expression of the gene can occur). A nucleic acid
molecule encoding the amino acid sequence of a B7-like
5 polypeptide may be amplified/expressed in prokaryotic,
yeast, insect (baculovirus systems), and/or eukaryotic
host cells. Selection of the host cell will depend in
part on whether a B7-like polypeptide is to be posttranslationally modified (e.g., glycosylated and/or
phosphorylated). If so, yeast, insect or mammalian
host cells are preferable. For a review of expression
vectors, see Meth. Enz., v.185, D.V. Goeddel, ed.
Academic Press Inc., San Diego, CA (1990).

Typically, expression vectors used in any of the 15 host cells will contain sequences for plasmid maintenance and for cloning and expression of exogenous nucleotide sequences. Such sequences, collectively referred to as "flanking sequences" in certain embodiments will typically include one or more of the 20 following nucleotide sequences: a promoter, one or more enhancer sequences, an origin οf replication, transcriptional termination sequence, a complete intron sequence containing a donor and acceptor splice site, a sequence encoding a leader sequence for polypeptide secretion, a ribosome binding site, a polyadenylation 25 sequence, a polylinker region for inserting the nucleic acid encoding the polypeptide to be expressed, and a selectable marker element. Each of these sequences is discussed below.

Optionally, the vector may contain a "tag"-encoding sequence, i.e., an oligonucleotide molecule located at the 5' or 3' end of a B7-like polypeptide

10

coding sequence; the oligonucleotide sequence encodes polyHis (such as hexaHis), or other "tag" such as FLAG, HA (hemaglutinin Influenza virus) or myc for which commercially available antibodies exist. This tag is typically fused to the polypeptide upon expression of the polypeptide, and can serve as a means for affinity purification of a B7-like polypeptide from the host cell. Affinity purification can be accomplished, for example, by column chromatography using antibodies against the tag as an affinity matrix. Optionally, the tag can subsequently be removed from a purified B7-like polypeptide by various means such as using certain peptidases for cleavage.

Flanking sequences may be homologous (i.e., from 15 the same species and/or strain as the host cell), heterologous (i.e., from a species other than the host cell species or strain), hybrid (i.e., a combination of flanking sequences from more than one source) synthetic, or the flanking sequences may be native 20 sequences which normally function to regulate B7-like polypeptide expression. As such, the source of a flanking sequence may be any prokaryotic or eukaryotic organism, any vertebrate or invertebrate organism, or any plant, provided that the flanking sequence 25 functional in, and can be activated by, the host cell machinery.

The flanking sequences useful in the vectors of this invention may be obtained by any of several methods well known in the art. Typically, flanking sequences useful herein other than B7-like gene flanking sequences will have been previously identified by mapping and/or by restriction endonuclease digestion

10

15

20

25

30

and can thus be isolated from the proper tissue source using the appropriate restriction endonucleases. In some cases, the full nucleotide sequence of a flanking sequence may be known. Here, the flanking sequence may be synthesized using the methods described herein for nucleic acid synthesis or cloning.

Where all or only a portion of the flanking sequence is known, it may be obtained using PCR and/or bу screening a genomic library with oligonucleotide and/or flanking sequence fragments from the same or another species. Where the flanking sequence is not known, a fragment of DNA containing a flanking sequence may be isolated from a larger piece of DNA that may contain, for example, a coding sequence or even another gene or genes. Isolation may be accomplished by restriction endonuclease digestion to produce the proper DNA fragment followed by isolation using agarose gel purification, Qiagen® column chromatography (Chatsworth, CA), or other methods known to the skilled artisan. The selection of suitable enzymes to accomplish this purpose will be readily apparent to one of ordinary skill in the art.

An origin of replication is typically a part of those prokaryotic expression vectors purchased commercially, and the origin aids in the amplification of the vector in a host cell. Amplification of the vector to a certain copy number can, in some cases, be important for the optimal expression of a B7-like polypeptide. If the vector of choice does not contain an origin of replication site, one may be chemically synthesized based on a known sequence and ligated into the vector. For example, the origin of replication from the plasmid pBR322 (Product No. 303-3s,

England Biolabs, Beverly, MA) is suitable for most Gram-negative bacteria and various origins (e.g., SV40, polyoma, adenovirus, vesicular stomatitus virus (VSV) or papillomaviruses such as HPV or BPV) are useful for cloning vectors in mammalian cells. Generally, the origin of replication component is not needed for mammalian expression vectors (for example, the SV40 origin is often used only because it contains the early promoter).

10 A transcription termination sequence is typically located 3' of the end of a polypeptide coding region and serves to terminate transcription. Usually, a transcription termination sequence in prokaryotic cells is a G-C rich fragment followed by a poly T sequence.

15 While the sequence is easily cloned from a library or even purchased commercially as part of a vector, it can also be readily synthesized using methods for nucleic acid synthesis such as those described herein.

A selectable marker gene element encodes a protein 20 necessary for the survival and growth of a host cell grown in a selective culture medium. Typical selection marker genes encode proteins that (a) confer resistance to antibiotics or other toxins, e.g., ampicillin, tetracycline, or kanamycin for prokaryotic host cells, 25 (b) complement auxotrophic deficiencies of the cell; or supply critical nutrients not available from complex media. Preferred selectable markers are the kanamycin resistance gene, the ampicillin resistance gene, and the tetracycline resistance gene. A neomycin 30 resistance gene may also be used for selection in prokaryotic and eukaryotic host cells.

10

15

20

Other selection genes may be used to amplify the gene which will be expressed. Amplification is the process wherein genes which are in greater demand for the production of a protein critical for growth are reiterated in tandem within the chromosomes successive generations of recombinant cells. Examples suitable selectable markers for mammalian cells include dihydrofolate reductase (DHFR) and thymidine kinase. The mammalian cell transformants are placed under selection pressure which only the transformants are uniquely adapted to survive by virtue of the selection gene present in the vector. pressure is imposed by culturing the transformed cells conditions in which the concentration selection agent in the medium is successively changed, thereby leading to the amplification of both the selection gene and the DNA that encodes a B7-like polypeptide. As a result, increased quantities of a B7-like polypeptide are synthesized from the amplified DNA.

A ribosome binding site is usually necessary for translation initiation of mRNA and is characterized by a Shine-Dalgarno sequence (prokaryotes) or a Kozak sequence (eukaryotes). The element is typically located 3' to the promoter and 5' to the coding sequence of a B7-like polypeptide to be expressed. The Shine-Dalgarno sequence is varied but is typically a polypurine (i.e., having a high A-G content). Many Shine-Dalgarno sequences have been identified, each of which can be readily synthesized using methods set forth herein and used in a prokaryotic vector.

A leader, or signal, sequence may be used to direct a B7-like polypeptide out of the host cell.

Typically, a nucleotide sequence encoding the signal sequence is positioned in the coding region of a B7like nucleic acid molecule, or directly at the 5' end of a B7-like polypeptide coding region. Many signal sequences have been identified, and any of those that are functional in the selected host cell may be used in conjunction with a B7-like nucleic acid molecule. Therefore, а signal sequence may be homologous (naturally occurring) or heterologous to a B7-like gene 10 Additionally, a signal sequence may be or cDNA. chemically synthesized using methods described herein. In most cases, the secretion of a B7-like polypeptide from the host cell via the presence of a signal peptide will result in the removal of the signal peptide from 15 the secreted B7-like polypeptide. The signal sequence may be a component of the vector, or it may be a part of a B7-like nucleic acid molecule that is inserted into the vector.

Included within the scope of this invention is the use of either a nucleotide sequence encoding a native 20 B7-like polypeptide signal sequence joined to a B7-like polypeptide coding region or a nucleotide sequence encoding a heterologous signal sequence joined to a B7like polypeptide coding region. The heterologous 25 signal sequence selected should be one that recognized and processed, i.e., cleaved by a signal peptidase, by the host cell. For prokaryotic host cells that do not recognize and process the native B7like polypeptide signal sequence, the signal sequence 30 substituted by a prokaryotic signal sequence selected, for example, from the group of the alkaline phosphatase, penicillinase, or heat-stable enterotoxin II leaders. For yeast secretion, the native B7-like

5

10

15

20

25

30

polypeptide signal sequence(s) may be substituted by the yeast invertase, alpha factor, or acid phosphatase leaders. In mammalian cell expression the native signal sequence is satisfactory, although other mammalian signal sequences may be suitable.

In some cases, such as where glycosylation is desired in a eukaryotic host cell expression system, one may manipulate the various presequences to improve glycosylation or yield. For example, one may alter the peptidase cleavage site of a particular signal peptide, or add presequences, which also may affect glycosylation. The final protein product may have, in the -1 position (relative to the first amino acid of the mature protein) one or more additional amino acids incident to expression, which may not have been totally For example, the final protein product may removed. have one or two amino acid residues found in the peptidase cleavage site, attached to the N-terminus. Alternatively, use of some enzyme cleavage sites may result in a slightly truncated form of the desired B7like polypeptide, if the enzyme cuts at such area within the mature polypeptide.

In many cases, transcription of a nucleic acid molecule is increased by the presence of one or more introns in the vector; this is particularly true where a polypeptide is produced in eukaryotic host cells, especially mammalian host cells. The introns used may naturally occurring be within а B7-like especially where the gene used is a full-length genomic sequence or a fragment thereof. Where the intron is not naturally occurring within the gene (as for most cDNAs), the intron(s) may be obtained from another The position of the intron with respect to source.

10

15

flanking sequences and a B7-like gene is generally important, as the intron must be transcribed to be effective. Thus, when a B7-like cDNA molecule is being transcribed, the preferred position for the intron is 3' to the transcription start site, and 5' to the polyA 5 transcription termination sequence. Preferably, the intron or introns will be located on one side or the other (i.e., 5' or 3') of the cDNA such that it does not interrupt the coding sequence. Any intron from any source, including any viral, prokaryotic and eukaryotic (plant or animal) organisms, may be used to practice this invention, provided that it is compatible with the host cell(s) into which it is inserted. Also included herein are synthetic introns. Optionally, more than one intron may be used in the vector.

The expression and cloning vectors of the present invention will each typically contain a promoter that is recognized by the host organism and operably linked to the molecule encoding a B7-like polypeptide. Promoters are untranscribed sequences located upstream 20 (5') to the start codon of a structural gene (generally within about 100 to 1000 bp) that control transcription of the structural gene. Promoters are conventionally grouped into one of two classes, 25 inducible promoters and constitutive promoters. Inducible promoters initiate increased levels transcription from DNA under their control in response to some change in culture conditions, such as the presence or absence of a nutrient or a change in 30 temperature. Constitutive promoters, on the other hand, initiate continual gene product production; that is, there is little or no control over gene expression. A large number of promoters, recognized by a variety of

5

10

25

30

potential host cells, are well known. A suitable promoter is operably linked to the DNA encoding a B7like polypeptide by removing the promoter from the source DNA by restriction enzyme digestion and inserting the desired promoter sequence into the vector. The native B7-like gene promoter sequence(s) may be used to direct amplification and/or expression of a B7-like nucleic acid molecule. A heterologous promoter is preferred, however, if it permits greater transcription and higher yields of the expressed protein as compared to the native promoter, and if it is compatible with the host cell system that has been selected for use.

Promoters suitable for use with prokaryotic hosts 15 include the beta-lactamase and lactose promoter alkaline phosphatase, a tryptophan promoter system; and hybrid promoters such as the tac promoter. Other known bacterial promoters are also suitable. Their sequences have been published, thereby 20 enabling one skilled in the art to ligate them to the desired DNA sequence(s), using linkers or adapters as needed to supply any useful restriction sites.

Suitable promoters for use with yeast hosts are also well known in the art. Yeast enhancers are advantageously used with yeast promoters. Suitable promoters for use with mammalian host cells are well known and include, but are not limited to, obtained from the genomes of viruses such as polyoma virus, fowlpox virus, adenovirus (such Adenovirus 2), bovine papilloma virus, avian sarcoma virus, cytomegalovirus (CMV), a retrovirus, hepatitis-B virus and most preferably Simian Virus 40

Other suitable mammalian promoters include heterologous mammalian promoters, e.g., heat-shock promoters and the actin promoter.

Additional promoters which may be of interest in controlling B7-like gene transcription include, but are limited to: the SV40 early promoter (Bernoist and Chambon, Nature, 290:304-310 (1981)); the CMV promoter; the promoter contained in the 3' long terminal repeat of Rous sarcoma virus (Yamamoto et al., 10 Cell, 22:787-797 (1980)); the herpes thymidine kinase promoter (Wagner et al., Proc. Natl. Acad. Sci. USA, 78:144-1445 (1981)); the regulatory sequences of the metallothionine gene (Brinster et al., Nature, 296:39-42 (1982)); prokaryotic expression vectors such as the 15 beta-lactamase promoter (Villa-Kamaroff, et al., Proc. Natl. Acad. Sci. USA, 75:3727-3731 (1978)); or the tac promoter (DeBoer, et al., Proc. Natl. Acad. Sci. USA, 80:21-25 (1983)). Also of interest are the following animal transcriptional control regions, which exhibit 20 tissue specificity and have been utilized in transgenic animals: the elastase I gene control region which is active in pancreatic acinar cells (Swift et al., Cell, 38:639-646 (1984); Ornitz et al., Cold Spring Harbor Quant. Biol., 50:399-409 (1986); MacDonald, 25 Hepatology, 7:425-515 (1987)); the insulin gene control region which is active in pancreatic beta (Hanahan, Nature, 315:115-122 (1985)); immunoglobulin gene control region which is active in lymphoid cells (Grosschedl et al., Cell, 38:647-658 30 (1984); Adames et al., Nature, 318:533-538 (1985); Alexander et al., Mol. Cell. Biol., 7:1436-1444 (1987)); the mouse mammary tumor virus control region

25

30

A-692

which is active in testicular, breast, lymphoid and mast cells (Leder et al., Cell, 45:485-495 (1986)); the albumin gene control region which is active in liver (Pinkert et al., Genes and Devel., 1:268-276 (1987)); the alphafetoprotein gene control region which 5 active in liver (Krumlauf et al., Mol. Cell. Biol., $\underline{5}$:1639-1648 (1985); Hammer et al., Science, $\underline{235}$:53-58 (1987))); the alpha 1-antitrypsin gene control region which is active in the liver (Kelsey et al., Genes and Devel., $\underline{1}$:161-171 (1987)); the beta-globin gene control 10 region which is active in myeloid cells (Mogram et al., Nature, 315:338-340 (1985); Kollias et al., Cell, $\underline{46}$:89-94 (1986)); the myelin basic protein gene control region which is active in oligodendrocyte cells in the brain (Readhead et al., Cell, 48:703-712 (1987)); the 15 myosin light chain-2 gene control region which is active in skeletal muscle (Sani, Nature, 314:283-286 (1985)); and the gonadotropic releasing hormone gene control region which is active in the hypothalamus 20 (Mason et al., Science, 234:1372-1378 (1986)).

An enhancer sequence may be inserted into the vector to increase the transcription of a DNA encoding an B7-like polypeptide of the present invention by higher eukaryotes. Enhancers are cis-acting elements of DNA, usually about 10-300 bp in length, that act on the promoter to increase transcription. Enhancers are relatively orientation and position independent. They have been found 5' and 3' to the transcription unit. Several enhancer sequences available from mammalian genes are known (e.g., globin, elastase, albumin, alpha-feto-protein and insulin). Typically, however, an enhancer from a virus will be used. The SV40 enhancer, the cytomegalovirus early promoter enhancer,

30

A-692

the polyoma enhancer, and adenovirus enhancers are exemplary enhancing elements for the activation of eukaryotic promoters. While an enhancer may be spliced into the vector at a position 5' or 3' to a B7-like nucleic acid molecule, it is typically located at a site 5' from the promoter.

Expression vectors of the invention may be constructed from a starting vector such as a commercially available vector. Such vectors may or may not contain all of the desired flanking sequences. Where one or more of the desired flanking sequences are not already present in the vector, they may be individually obtained and ligated into the vector. Methods used for obtaining each of the flanking sequences are well known to one skilled in the art.

Preferred vectors for practicing this invention are those which are compatible with bacterial, insect, and mammalian host cells. Such vectors include, inter alia, pCRII, pCR3, and pcDNA3.1 (Invitrogen Company, Carlsbad, CA), pBSII (Stratagene Company, La Jolla, CA), pET15. (Novagen, Madison, WI), pGEX (Pharmacia Biotech, Piscataway, NJ), pEGFP-N2 (Clontech, Palo Alto, CA), pETL (BlueBacII; Invitrogen), pDSR-alpha (PCT Publication No. W090/14363) and pFastBacDual (Gibco/BRL, Grand Island, NY).

Additional suitable vectors include, but are not limited to, cosmids, plasmids or modified viruses, but it will be appreciated that the vector system must be compatible with the selected host cell. Such vectors include, but are not limited to plasmids such as Bluescript plasmid derivatives (a high copy number ColE1-based phagemid, Stratagene Cloning Systems Inc.,

10

15

20

La Jolla CA), PCR cloning plasmids designed for cloning Taq-amplified PCR products (e.g., TOPOTM TA Cloning Kit, PCR2.1 plasmid derivatives, Invitrogen, Carlsbad, CA), and mammalian, yeast, or virus vectors such as a baculovirus expression system (pBacPAK plasmid derivatives, Clontech, Palo Alto, CA).

After the vector has been constructed and nucleic acid molecule encoding a B7-like polypeptide has been inserted into the proper site of the vector, the completed vector may be inserted into a suitable host cell for amplification and/or polypeptide expression. The transformation of an expression vector for a B7-like polypeptide into a selected host cell may be accomplished by well known methods including methods such as transfection, infection, calcium chloride, electroporation, microinjection, lipofection or DEAE-dextran method or other known techniques. method selected will in part be a function of the type of host cell to be used. These methods and other suitable methods are well known to the skilled artisan, and are set forth, for example, in Sambrook et al., supra.

Host cells may be prokaryotic host cells (such as E. coli) or eukaryotic host cells (such as a yeast 25 cell, an insect cell or a vertebrate cell). The host cell, when cultured under appropriate conditions, synthesizes а B7-like polypeptide which subsequently be collected from the culture medium (if the host cell secretes it into the medium) or directly 30 from the host cell producing it (if it is secreted). The selection of an appropriate host cell will depend upon various factors, such as desired expression levels, polypeptide modifications that are

desirable or necessary for activity, such as glycosylation or phosphorylation, and ease of folding into a biologically active molecule.

A number of suitable host cells are known in the and many are available from the American Type Culture Collection (ATCC), 10801 University Boulevard, Manassas, VA 20110-2209. Examples include, but are not limited to, mammalian cells, such as Chinese hamster ovary cells (CHO) (ATCC No. CCL61) CHO DHFR-10 cells (Urlaub et al., Proc. Natl. Acad. Sci. 97:4216-4220 (1980)), human embryonic kidney (HEK) 293 or 293T cells (ATCC No. CRL1573), or 3T3 cells (ATCC No. CCL92). The selection of suitable mammalian host cells methods and for transformation, culture. 15 amplification, screening and product production and purification are known in the art. Other suitable mammalian cell lines, are the monkey COS-1 (ATCC No. CRL1650) and COS-7 cell lines (ATCC No. CRL1651), and the CV-1 cell line (ATCC No. CCL70). Further exemplary 20 mammalian host cells include primate cell lines and rodent cell lines, including transformed cell lines. Normal diploid cells, cell strains derived from in vitro culture of primary tissue, as well as primary explants, are also suitable. Candidate cells may be 25 genotypically deficient in the selection gene, or may contain a dominantly acting selection gene. Other suitable mammalian cell lines include but are not limited to, mouse neuroblastoma N2A cells, HeLa, mouse L-929 cells, 3T3 lines derived from Swiss, Balb-c or 30 NIH mice, BHK or HaK hamster cell lines, which are available from the ATCC. Each of these cell lines is known by and available to those skilled in the art of protein expression.

25

30

A-692

Similarly useful as host cells suitable for the present invention are bacterial cells. For example, the various strains of $E.\ coli\ (e.g.,\ HB101,\ (ATCC\ No.\ 33694)\ DH5\alpha,\ DH10,\ and\ MC1061\ (ATCC\ No.\ 53338))$ are well-known as host cells in the field of biotechnology. Various strains of $B.\ subtilis,\ Pseudomonas\ spp.$, other $Bacillus\ spp.$, $Streptomyces\ spp.$, and the like may also be employed in this method.

Many strains of yeast cells known to those skilled in the art are also available as host cells for the expression of the polypeptides of the present invention. Preferred yeast cells include, for example, Saccharomyces cerivisae and Pichia pastoris.

Additionally, where desired, insect cell systems 15 may be utilized in the methods of the present invention. Such systems are described for example in Kitts et al., Biotechniques, 14:810-817 (1993);Lucklow, Curr. Opin. Biotechnol., $\underline{4}:564-572$ (1993); and Lucklow et al. *Virol.*, 67:4566-4579 (J.(1993).Preferred insect cells are Sf-9 and Hi5 (Invitrogen, 20 Carlsbad, CA).

One may also use transgenic animals to express glycosylated B7-like polypeptides. For example, one may use a transgenic milk-producing animal (a cow or goat, for example) and obtain the present glycosylated polypeptide in the animal milk. One may also use plants to produce B7-like polypeptides, however, in general, the glycosylation occurring in plants is different from that produced in mammalian cells, and may result in a glycosylated product which is not suitable for human therapeutic use.

10

15

20

25

Polypeptide Production

Host cells comprising a B7-like polypeptide expression vector may be cultured using standard media well known to the skilled artisan. The media will usually contain all nutrients necessary for the growth survival of the cells. Suitable media culturing E. coli cells include, for example, Luria Broth (LB) and/or Terrific Broth (TB). Suitable media for culturing eukaryotic cells include Roswell Park Memorial Institute medium 1640 (RPMI 1640), Minimal Essential Medium (MEM) and/or Dulbecco's Modified Eagle Medium (DMEM), all of which may be supplemented with and/or growth factors as indicated by the particular cell line being cultured. A suitable medium for insect cultures is Grace's medium supplemented with yeastolate, lactalbumin hydrolysate and/or fetal calf serum, as necessary.

Typically, an antibiotic or other compound useful for selective growth of transformed cells is added as a supplement to the media. The compound to be used will be dictated by the selectable marker element present on the plasmid with which the host cell was transformed. For example, where the selectable marker element is kanamycin resistance, the compound added to the culture medium will be kanamycin. Other compounds for selective growth include ampicillin, tetracycline, and neomycin.

The amount of a B7-like polypeptide produced by a host cell can be evaluated using standard methods known in the art. Such methods include, without limitation, Western blot analysis, SDS-polyacrylamide gel electrophoresis, non-denaturing gel electrophoresis,

HPLC separation, immunoprecipitation, and/or activity assays such as DNA binding gel shift assays.

If a B7-like polypeptide has been designed to be secreted from the host cells, the majority of polypeptide may be found in the cell culture medium. If however, a B7-like polypeptide is not secreted from the host cells, it will be present in the cytoplasm and/or the nucleus (for eukaryotic host cells) or in the cytosol (for bacterial host cells).

For a B7-like polypeptide situated in the host cell cytoplasm and/or the nucleus (for eukaryotic host cells) or in the cytosol (for bacterial host cells), intracellular material (including inclusion bodies for gram-negative bacteria) can be extracted from the host cell using any standard technique known to the skilled artisan. For example, the host cells can be lysed to release the contents of the periplasm/cytoplasm by French press, homogenization, and/or sonication followed by centrifugation.

20 a B7-like polypeptide has formed inclusion bodies in the cytosol, the inclusion bodies can often bind to the inner and/or outer cellular membranes and thus will be found primarily in the pellet material after centrifugation. The pellet material can then be 25 treated at pH extremes or with a chaotropic agent such as a detergent, guanidine, guanidine derivatives, urea, or urea derivatives in the presence of a reducing agent such as dithiothreitol at alkaline pH or carboxyethyl phosphine at acid pH to release, break 30 apart, and solubilize the inclusion bodies. A B7-like polypeptide in its now soluble form can analyzed using gel electrophoresis, immunoprecipitation

or the like. If it is desired to isolate a B7-like polypeptide, isolation may be accomplished using standard methods such as those described herein and in Marston et al., Meth. Enz., $\underline{182}:264-275$ (1990).

5 In some cases, a B7-like polypeptide may not be biologically active upon isolation. Various methods for "refolding" or converting the polypeptide to its tertiary structure and generating disulfide linkages can be used to restore biological activity. methods include exposing the solubilized polypeptide to 10 a pH usually above 7 and in the presence of a particular concentration of a chaotrope. The selection of chaotrope is very similar to the choices used for inclusion body solubilization, but usually 15 chaotrope is used at a lower concentration and is not necessarily the same as chaotropes used for solubilization. In most cases the refolding/oxidation solution will also contain a reducing agent or the reducing agent plus its oxidized form in a specific ratio to generate a particular redox potential allowing 20 for disulfide shuffling to occur in the formation of the protein's cysteine bridge(s). Some of the commonly used redox couples include cysteine/cystamine, glutathione (GSH)/dithiobis GSH, cupric chloride, 25 dithiothreitol(DTT)/ dithiane DTT, and 2mercaptoethanol(bME)/dithio-b(ME). A cosolvent may be used to increase the efficiency of the refolding, and the more common reagents used for this purpose include glycerol, polyethylene glycol of various molecular 30 weights, arginine and the like.

Ιf inclusion bodies are not formed to а significant degree upon expression of а B7-like polypeptide, then the polypeptide will found be

20

30

A-692

primarily in the supernatant after centrifugation of the cell homogenate. The polypeptide may be further isolated from the supernatant using methods such as those described herein.

The purification of a B7-like polypeptide from solution can be accomplished using a variety of techniques. If the polypeptide has been synthesized such that it contains a tag such as Hexahistidine (B7-like polypeptide/hexaHis) or another small peptide such as FLAG (Eastman Kodak Co., New Haven, CT) or myc (Invitrogen, Carlsbad, CA) at either its carboxyl or amino terminus, it may be purified in a one-step process by passing the solution through an affinity column where the column matrix has a high affinity for the tag.

For example, polyhistidine binds with great affinity and specificity to nickel, thus an affinity column of nickel (such as the Qiagen® nickel columns) can be used for purification of a B7-like polypeptide/polyHis. See for example, Ausubel et al., eds., Current Protocols in Molecular Biology, Section 10.11.8, John Wiley & Sons, New York (1993).

Additionally, a B7-like polypeptide may be purified through the use of a monoclonal antibody which is capable of specifically recognizing and binding to a B7-like polypeptide.

Suitable procedures for purification thus include, without limitation, affinity chromatography, immunoaffinity chromatography, ion exchange chromatography, molecular sieve chromatography, High Performance Liquid Chromatography (HPLC),

5

electrophoresis (including native gel electrophoresis) followed by gel elution, and preparative isoelectric focusing ("Isoprime" machine/technique, Hoefer Scientific, San Francisco, CA). In some cases, two or more purification techniques may be combined to achieve increased purity.

B7-like polypeptides may also be prepared by chemical synthesis methods (such as solid phase peptide synthesis) using techniques known in the art, such as 10 those set forth by Merrifield et al., J. Am. Chem. Soc., 85:2149 (1963), Houghten et al., Proc Natl Acad. Sci. USA, 82:5132 (1985), and Stewart and Young, Solid Phase Peptide Synthesis, Pierce Chemical Co., Rockford, IL (1984). Such polypeptides may be synthesized with 15 without a methionine on the amino terminus. Chemically synthesized B7-like polypeptides may be oxidized using methods set forth in these references to form disulfide bridges. Chemically synthesized B7-like polypeptides are expected to have comparable biological 20 activity to the corresponding B7-like polypeptides recombinantly or purified from produced sources, and thus may be used interchangeably with a recombinant or natural B7-like polypeptide.

Another means of obtaining a B7-like polypeptide is via purification from biological samples such as 25 source tissues and/or fluids in which a B7-like polypeptide is naturally found. Such purification can be conducted using methods for protein purification as described herein. The presence οf a 30 polypeptide during purification may be monitored using, for example, an antibody prepared against recombinantly produced B7-like polypeptide or peptide

fragments thereof.

A number of additional methods for producing nucleic acids and polypeptides are known in the art, used to produce polypeptides and can be specificity for a B7-like polypeptide. See for example, Roberts et al., Proc. Natl. Acad. Sci., 94:12297-12303 which describes the production of proteins between an mRNA and its encoded peptide. also Roberts, R., Curr. Opin. Chem. Biol., 3:268-273 10 (1999).Additionally, U.S. Patent No. 5,824,469 describes methods of obtaining oligonucleotides capable of carrying out a specific biological function. procedure involves generating a heterogeneous pool of oligonucleotides, each having a 5' randomized sequence, 15 a central preselected sequence, and a 3' randomized sequence. The resulting heterogeneous pool introduced into a population of cells that do not exhibit the desired biological function. Subpopulations of the cells are then screened for those which exhibit a predetermined biological 20 function. From that subpopulation, oligonucleotides capable of carrying out the desired biological function are isolated.

U.S. Patent Nos. 5,723,323, 5,763,192, 5,814,476
and 5,817,483 describe processes for producing peptides
or polypeptides. This is done by producing stochastic
genes or fragments thereof, and then introducing these
genes into host cells which produce one or more
proteins encoded by the stochastic genes. The host
cells are then screened to identify those clones
producing peptides or polypeptides having the desired
activity.

10

15

20

25

30

Chemical Derivatives

Chemically modified derivatives of B7-like polypeptides may be prepared by one skilled in the art, given the disclosures set forth hereinbelow. polypeptide derivatives are modified in a manner that is different, either in the type or location of the molecules naturally attached to the polypeptide. Derivatives include molecules may formed by deletion of one or more naturally-attached chemical groups. A polypeptide comprising an amino sequence of SEQ ID NOs: 2, 4, 6, 8, 10, 12 or 14, or a B7-like polypeptide variant may be modified by the covalent attachment of one or more polymers. For example, the polymer selected is typically water soluble so that the protein to which it is attached does not precipitate in an aqueous environment, such as a physiological environment. Included within the scope suitable polymers is a mixture of polymers. Preferably, for therapeutic use of the end-product preparation, the polymer will be pharmaceutically acceptable.

The polymers each may be of any molecular weight and may be branched or unbranched. The polymers each typically have an average molecular weight of between about 2kDa to about 100kDa (the term "about" indicating that in preparations of a water soluble polymer, some molecules will weigh more, some less, than the stated molecular weight). The average molecular weight of each polymer preferably is between about 5kDa and about 50kDa, more preferably between about 12kDa and about 40kDa and most preferably between about 20kDa and about 35kDa.

10

15

Suitable water soluble polymers or mixtures thereof include, but are not limited to, N-linked or Olinked carbohydrates, sugars, phosphates, polyethylene glycol (PEG) (including the forms of PEG that have been used to derivatize proteins, including mono- (C_1-C_{10}) alkoxy- or aryloxy-polyethylene glycol), monomethoxypolyethylene glycol, dextran (such as low molecular weight dextran, of, for example about 6 kD), cellulose, or other carbohydrate based polymers, poly-(N-vinyl pyrrolidone) polyethylene glycol, propylene glycol homopolymers, a polypropylene oxide/ethylene oxide copolymer, polyoxyethylated polyols (e.g., glycerol) and polyvinyl alcohol. Also encompassed by the present invention are bifunctional crosslinking molecules which may be used to prepare covalently attached multimers of a polypeptide comprising an amino acid sequence of SEQ ID NOs: 2, 4, 6, 8, 10, 12 or 14 or a B7-like polypeptide variant.

In general, chemical derivatization may 20 performed under any suitable condition used to react a protein with an activated polymer molecule. for preparing chemical derivatives of polypeptides will generally comprise the steps of (a) reacting polypeptide with the activated polymer molecule (such 25 as a reactive ester or aldehyde derivative of the polymer molecule) under conditions whereby polypeptide comprising the amino acid sequence of SEQ ID NOs: 2, 4, 6, 8, 10, 12 or 14, or a B7-like polypeptide variant becomes attached to one or more 30 polymer molecules, and (b) obtaining the reaction The optimal reaction conditions will be product(s). determined based on known parameters and the desired result. For example, the larger the ratio of polymer

5

molecules:protein, the greater the percentage of attached polymer molecule. In one embodiment, a B7-like polypeptide derivative may have a single polymer molecule moiety at the amino terminus. See, for example, U.S. Patent No. 5,234,784.

The pegylation of the polypeptide specifically may be carried out by any of the pegylation reactions known in the art, as described for example in the following references: Francis et al., Focus on Growth Factors, 10 3:4-10(1992);EP 0154316; EP 0401384 and U.S. Patent No. 4,179,337. For example, pegylation may be carried out via an acylation reaction or an alkylation reaction with a reactive polyethylene glycol molecule (or an analogous reactive water-soluble polymer) as described herein. 15 For the acylation reactions, the polymer(s) selected should have a single reactive ester group. reductive alkylation, the polymer(s) selected should have a single reactive aldehyde group. A reactive 20 aldehyde is, for example, polyethylene propional dehyde, which is water stable, or mono $C_1\text{-}C_{10}$ alkoxy or aryloxy derivatives thereof (see U.S. Patent No. 5,252,714).

In another embodiment, B7-like polypeptides may be chemically coupled to biotin, and the biotin/B7-like 25 polypeptide molecules which are conjugated are then allowed to bind to avidin, resulting in tetravalent avidin/biotin/B7-like polypeptide molecules. B7-like polypeptides may also be covalently coupled 30 dinitrophenol (DNP) or trinitrophenol (TNP) and the resulting conjugates precipitated with anti-DNP

10

25

30

anti-TNP-IgM to form decameric conjugates with a valency of 10.

Generally, conditions which may be alleviated or modulated by the administration of the present B7-like polypeptide derivatives include those described herein for B7-like polypeptides. However, the B7-like polypeptide derivatives disclosed herein may have additional activities, enhanced or reduced biological activity, or other characteristics, such as increased or decreased half-life, as compared to the non-derivatized molecules.

Genetically Engineered Non-Human Animals

Additionally included within the scope of the present invention are non-human animals such as mice, rats, or other rodents, rabbits, goats, or sheep, or other farm animals, in which the genes encoding the native B7-like polypeptides have been disrupted ("knocked out") such that the level of expression of this gene or genes is (are) significantly decreased or completely abolished. Such animals may be prepared using techniques and methods such as those described in U.S. Patent No. 5,557,032.

The present invention further includes non-human animals such as mice, rats, or other rodents, rabbits, goats, sheep, or other farm animals, in which either the native form of the B7-like genes for that animal or a heterologous B7-like genes are over-expressed by the animal, thereby creating a "transgenic" animal. Such transgenic animals may be prepared using well known methods such as those described in U.S. Patent No 5,489,743 and PCT application No. W094/28122.

5

10

15

20

25

The present invention further includes non-human animals in which the promoter for one or more of the B7-like polypeptides of the present invention is either activated or inactivated (e.g., by using homologous recombination methods) to alter the level of expression of one or more of the native B7-like polypeptides.

These non-human animals may be used for drug candidate screening. In such screening, the impact of a drug candidate on the animal may be measured. example, drug candidates may decrease or increase the expression of a B7-like gene. In certain embodiments, the amount of a B7-like polypeptide that is produced may be measured after the exposure of the animal to the drug candidate. Additionally, in certain embodiments, one may detect the actual impact of the drug candidate on the animal. For example, the overexpression of a particular gene may result in, or be associated with, a disease or pathological condition. In such cases, one may test a drug candidate's ability to decrease expression of the gene or its ability to prevent or inhibit a pathological condition. In other examples, the production of a particular metabolic product such as a fragment of a polypeptide, may result in, or be associated with, a disease or pathological condition. In such cases, one may test a drug candidate's ability to decrease the production of such a metabolic product or its ability to prevent or inhibit a pathological condition.

30 Microarray

It will be appreciated that DNA microarray technology can be utilized in accordance with the present invention. DNA microarrays are miniature, high

10

15

20

25

density arrays of nucleic acids positioned on a solid support, such as glass. Each cell or element within the array has numerous copies of a single species of DNA which acts as a target for hybridization for its cognate mRNA. In expression profiling using DNA microarray technology, mRNA is first extracted from a cell or tissue sample and then converted enzymatically to fluorescently labeled cDNA. This material hybridized to the microarray and unbound cDNA is removed by washing. The expression of discrete genes represented on the array is then visualized quantitating the amount of labeled cDNA which specifically bound to each target DNA. In this way, the expression of thousands of genes can be quantitated in a high throughput, parallel manner from a single sample of biological material.

This high throughput expression profiling has a broad range of applications with respect to the B7-like molecules of the invention, including, but not limited to: the identification and validation of B7-like genes in disease and as targets for therapeutics; molecular toxicology of B7-like molecules and inhibitors thereof; stratification of populations and generation surrogate markers for clinical trials; and enhancing B7-like small molecule drug discovery by aiding in the identification of selective compounds in high throughput screens (HTS).

Selective Binding Agents

As used herein, the term "selective binding agent" refers to a molecule which has specificity for one or more B7-like polypeptides. Suitable selective binding

regions.

30

agents include, but are not limited to, antibodies and derivatives thereof, polypeptides, and small molecules. Suitable selective binding agents may be prepared using methods known in the art. An exemplary B7-like polypeptide selective binding agent of the present invention is capable of binding a certain portion of at least one B7-like polypeptide, thereby inhibiting the binding of such polypeptide to the B7-like polypeptide receptor(s).

10 Selective binding agents such as antibodies and antibody fragments that each bind at least one B7-like polypeptide are within the scope of the present invention. The antibodies may be polyclonal including monospecific polyclonal, monoclonal (MAbs), recombinant, chimeric, humanized such as CDR-grafted, 15 human, single chain, and/or bispecific, as well as fragments, variants or derivatives thereof. fragments include those portions of the antibody which bind to an epitope on at least one B7-like polypeptide. 20 Examples of such fragments include Fab and F(ab') fragments generated by enzymatic cleavage of fulllength antibodies. Other binding fragments include those generated by recombinant DNA techniques, such as expression of recombinant plasmids containing 25 nucleic acid sequences encoding antibody variable

Polyclonal antibodies directed toward at least one B7-like polypeptide generally are produced in animals (e.g., rabbits or mice) by means of multiple subcutaneous or intraperitoneal injections of a B7-like polypeptide and an adjuvant. It may be useful to conjugate a B7-like polypeptide to a carrier protein

5

25

30

that is immunogenic in the species to be immunized, such as keyhole limpet heocyanin, serum, albumin, bovine thyroglobulin, or soybean trypsin inhibitor. Also, aggregating agents such as alum are used to enhance the immune response. After immunization, the animals are bled and the serum is assayed for anti-B7-like polypeptide antibody titer.

Monoclonal antibodies directed toward at least one B7-like polypeptide are produced using any method which provides for the production of antibody molecules by 10 continuous cell lines in culture. Examples of suitable methods for preparing monoclonal antibodies include the hybridoma methods of Kohler et al., Nature, 256:495-497 (1975) and the human B-cell hybridoma method, Kozbor, 15 J. Immunol., 133:3001 (1984);Brodeur al., Monoclonal Antibody Production Techniques and Applications, pp. 51-63 (Marcel Dekker, Inc., New York (1987)). Also provided by the invention are hybridoma cell lines which produce monoclonal antibodies reactive 20 with at least one B7-like polypeptide.

Monoclonal antibodies of the invention may be modified for use as therapeutics. One embodiment is a "chimeric" antibody in which a portion of the heavy and/or light chain is identical with or homologous to a corresponding sequence in antibodies derived from a particular species or belonging to a antibody class or subclass, while the remainder of the chain(s) is identical with or homologous corresponding sequence in antibodies derived another species or belonging to another antibody class or subclass. Also included are fragments of such antibodies, so long as they exhibit the desired

20

25

30

A-692

biological activity. See, U.S. Patent No. 4,816,567; Morrison *et al.*, *Proc. Natl. Acad. Sci.*, <u>81</u>:6851-6855 (1985).

In another embodiment, a monoclonal antibody of 5 the invention is a "humanized" antibody. Methods for humanizing non-human antibodies are well known in the See U.S. Patent Nos. 5,585,089, and 5,693,762. Generally, a humanized antibody has one or more amino acid residues introduced into it from a source which is 10 non-human. Humanization can be performed, for example, using methods described in the art (Jones et al., Nature, 321:522-525 (1986); Riechmann et al., Nature, 332:323-327 (1988); Verhoeyen et al., Science, 239:1534-1536 (1988)), by substituting at least a portion of a rodent complementarity-determining region 15 the corresponding regions (CDR) for of а human antibody.

Also encompassed by the invention are human antibodies which bind at least one B7-like polypeptide. Using transgenic animals (e.g., mice) that are capable of producing a repertoire of human antibodies in the absence of endogenous immunoglobulin production, such antibodies are produced by immunization with a B7-like antigen (i.e., having at least 6 contiguous amino acids), optionally conjugated to a carrier. See, for example, Jakobovits et al., Proc. Natl. Acad. Sci., 90:2551-2555 (1993); Jakobovits et al., Nature <u>362</u>:255-258 (1993); Bruggermann et al., Year in Immuno., 7:33 (1993).In one method, such transgenic animals are produced by incapacitating the endogenous loci encoding the heavy and light immunoglobulin chains therein, and inserting loci encoding human heavy and light chain

10

15

proteins into the genome thereof. Partially modified animals, that is those having less than the full complement of modifications, are then cross-bred to obtain an animal having all of the desired immune system modifications. When administered an immunogen, these transgenic animals produce antibodies with human (rather than e.g., murine) amino acid sequences, including variable regions which are immunospecific for these antigens. See PCT Application PCT/US96/05928 and PCT/US93/06926. Additional methods described in U.S. Patent No. 5,545,807, Application NOs. PCT/US91/245 and PCT/GB89/01207, and in EP 546073B1 and EP 546073A1. Human antibodies may also be produced by the expression of recombinant DNA in host cells or by expression in hybridoma cells as described herein.

In an alternative embodiment, human antibodies can be produced from phage-display libraries (Hoogenboom et al., J. Mol. Biol., 227:381 (1991); Marks et al., J. 20 Mol. Biol., 222:581 (1991)). These processes mimic selection through the display immune of antibody repertoires on the surface of filamentous bacteriophage, and subsequent selection of phage by their binding to an antigen of choice. One such 25 technique is described in PCTApplication PCT/US98/17364, which describes the isolation of high affinity and functional agonistic antibodies for MPLand msk- receptors using such an approach.

Chimeric, CDR grafted, and humanized antibodies are typically produced by recombinant methods. Nucleic acids encoding the antibodies are introduced into host cells and expressed using materials and procedures

5

10

15

20

25

30

described herein. In a preferred embodiment, the antibodies are produced in mammalian host cells, such as CHO cells. Monoclonal (e.g., human) antibodies may be produced by the expression of recombinant DNA in host cells or by expression in hybridoma cells as described herein.

The anti-B7-like antibodies of the invention may be employed in any known assay method, such as competitive binding assays, direct and indirect sandwich assays, and immunoprecipitation assays (Sola, Monoclonal Antibodies: A Manual of Techniques, pp. 147-158, CRC Press, Inc. (1987)) for the detection and quantitation of B7-like polypeptides. The antibodies will bind B7-like polypeptides with an affinity which is appropriate for the assay method being employed.

For diagnostic applications, certain in embodiments, anti-B7-like antibodies may be labeled with a detectable moiety. The detectable moiety can be any one which is capable of producing, either directly or indirectly, a detectable signal. For example, the detectable moiety may be a radioisotope, such as ³H, $^{14}\mathrm{C}$, $^{32}\mathrm{P}$, $^{35}\mathrm{S}$, or $^{125}\mathrm{I}$, a fluorescent or chemiluminescent compound, such as fluorescein isothiocyanate, rhodamine, or luciferin; or an enzyme, such as alkaline phosphatase, β -galactosidase, or horseradish peroxidase (Bayer et al., Meth. Enz., <u>184</u>:138-163 (1990)).

Competitive binding assays rely on the ability of a labeled standard (e.g., a B7-like polypeptide, or an immunologically reactive portion thereof) to compete with the test sample analyte (a B7-like polypeptide) for binding with a limited amount of anti-B7-like

30

A-692

antibody. The amount of B7-like polypeptide in the test sample is inversely proportional to the amount of standard that becomes bound to the antibodies. То facilitate determining the amount of standard that becomes bound, the antibodies typically insolubilized before or after the competition, so that standard and analyte that are bound the antibodies may conveniently be separated from the standard and analyte which remain unbound.

10 Sandwich assays typically involve the use of two antibodies, each capable of binding to a different immunogenic portion, or epitope, of the protein to be detected and/or quantitated. In a sandwich assay, the test sample analyte is typically bound by a first antibody which is immobilized on a solid support, and 15 thereafter a second antibody binds to the analyte, thus forming an insoluble three part complex. See, e.g., U.S. Patent No. 4,376,110. The second antibody may itself be labeled with a detectable moiety (direct 20 sandwich assays) or may be measured using an antiimmunoglobulin antibody that labeled is with detectable moiety (indirect sandwich assays). For example, one type of sandwich assay is an enzyme-linked immunosorbent assay (ELISA), in which case the 25 detectable moiety is an enzyme.

The selective binding agents, including anti-B7like antibodies, also are useful for in vivo imaging. An antibody labeled with a detectable moiety may be administered to an animal, preferably into the bloodstream, and the presence and location of labeled antibody in the host is assayed. The antibody may be labeled with any moiety that is detectable in an

10

30

magnetic resonance, animal, whether by nuclear radiology, or other detection means known in the art.

the invention, agents οf binding Selective including antibodies, may be used as therapeutics. These therapeutic agents are generally agonists or antagonists, in that they either enhance or reduce, respectively, at least one of the biological activities polypeptide. one B7-like least embodiment, antagonist antibodies of the invention are antibodies or binding fragments thereof which are capable of specifically binding to at least one B7-like polypeptide and which are capable of inhibiting or eliminating the functional activity of the mature form of at least one B7-like polypeptide in vivo or in vitro. In preferred embodiments, the selective binding 15 agent, e.g., an antagonist antibody, will inhibit the functional activity of the mature form of at least one 50%, about B7-like polypeptide by at least In another least about 80%. preferably by at embodiment, the selective binding agent may be an 20 antibody that is capable of interacting with a B7-like receptor) thereby ligand or binding partner (a inhibiting or eliminating B7-like activity in vitro or Selective binding agents, including agonist in vivo. and antagonist anti-B7-like antibodies, are identified 25 by screening assays which are well known in the art.

The invention also relates to a kit comprising B7like selective binding agents (such as antibodies) and other reagents useful for detecting B7-like polypeptide Such reagents may levels in biological samples. include, a detectable label, blocking serum, positive and negative control samples, and detection reagents.

5

10

15

20

25

30

The B7-like polypeptides of the present invention can be used to clone B7-like receptors, using an expression cloning strategy. Radiolabeled (125-Iodine) B7-like polypeptide or affinity/activity-tagged B7-like polypeptide (such as an Fc fusion or an alkaline phosphatase fusion) can be used in binding assays to identify a cell type or cell line or tissue that expresses B7-like receptor(s). RNA isolated from such cells or tissues can be converted to cDNA, cloned into a mammalian expression vector, and transfected into mammalian cells (such as COS or 293 cells) to create an expression library. A radiolabeled or tagged B7-like polypeptide can then be used as an affinity ligand to identify and isolate from this library the subset of cells which express the B7-like receptor(s) on their surface. DNA can then be isolated from these cells and transfected into mammalian cells to create a secondary expression library in which the fraction of cells expressing B7-like receptor(s) is many-fold higher than in the original library. This enrichment process can be repeated iteratively until a single recombinant clone containing a B7-like receptor is isolated. Isolation of the B7-like receptor(s) is useful for identifying developing or novel agonists antagonists of B7-like polypeptide signaling the pathway. Such agonists and antagonists include soluble B7-like receptor(s), anti-B7-like receptor antibodies, small molecules, or antisense oligonucleotides, and they for treating, preventing, may be used diagnosing one or more disease or disorders, including those described herein.

5

10

Assaying for other modulators of B7-like polypeptide activity

In some situations, it may be desirable identify molecules that are modulators, i.e., agonists or antagonists, of the activity of "the mature form of at least one B7-like polypeptide. Natural or synthetic molecules that modulate at least one B7-like polypeptide may be identified using one screening assays, such as those described herein. Such molecules may be administered either in an ex vivo manner, or in an in vivo manner by injection, or by oral delivery, implantation device, or the like.

"Test molecule(s)" refers to the molecule(s) that is/are under evaluation for the ability to modulate (i.e., increase or decrease) the activity of 15 mature form of at least one B7-like polypeptide. commonly, a test molecule will interact directly with the mature form of at least one B7-like polypeptide. However, it is also contemplated that a test molecule 20 may also modulate B7-like polypeptide activity indirectly, such as by affecting B7-like expression, or by binding to a B7-like binding partner (e.g., receptor or ligand). In one embodiment, a test molecule will bind to at least one B7-like polypeptide with an affinity constant of at least about 10^{-6} M, 25 preferably about 10^{-8} M, more preferably about 10^{-9} M, and even more preferably about 10^{-10} M.

Methods for identifying compounds which interact with at least one B7-like polypeptide are encompassed by the present invention. In certain embodiments, a B7-like polypeptide is incubated with a test molecule

5

10

15

20

25

under conditions which permit the interaction of the test molecule with a B7-like polypeptide, and the extent of the interaction can be measured. The test molecule(s) can be screened in a substantially purified form or in a crude mixture.

In certain embodiments, a B7-like polypeptide agonist or antagonist may be a protein, peptide, carbohydrate, lipid, or small molecular weight molecule which interacts with at least one B7-like polypeptide to regulate its activity. Molecules which regulate B7-like polypeptide expression include nucleic acids which are complementary to nucleic acids encoding a B7-like polypeptide, or are complementary to nucleic acids sequences which direct or control the expression of at least one B7-like polypeptide, and which act as antisense regulators of expression.

Once a set of test molecules has been identified as interacting with at least one B7-like polypeptide, the molecules may be further evaluated for their ability to increase or decrease B7-like polypeptide activity. The measurement of the interaction of test molecules with at least one B7-like polypeptide may be carried out in several formats, including cell-based binding assays, membrane binding assays, solution-phase assays and immunoassays. In general, test molecules are incubated with a B7-like polypeptide specified period of time, and B7-like polypeptide activity is determined by one or more assays for measuring biological activity.

The interaction of test molecules with at least one B7-like polypeptide may also be assayed directly

using polyclonal or monoclonal antibodies in an immunoassay. Alternatively, modified forms of at least one B7-like polypeptide containing epitope tags as described herein may be used in immunoassays.

5 In the event that B7-like polypeptides display biological activity through an interaction with a binding partner (e.g., a receptor or a ligand), a variety of in vitro assays may be used to measure the binding of such B7-like polypeptides 10 corresponding binding partner (such as a selective binding agent, receptor or ligand). These assays may be used to screen test molecules for their ability to increase or decrease the rate and/or the extent of binding of a B7-like polypeptide to its binding 15 partner. In one assay, a B7-like polypeptide is immobilized in the wells of a microtiter plate. Radiolabeled B7-like binding partner (for example, iodinated B7-like binding partner) and the test molecule(s) can then be added either one at a time (in either order) or simultaneously to the wells. 20 incubation, the wells can be washed and counted, using a scintillation counter, for radioactivity to determine the extent to which the binding partner bound to the B7-like polypeptide. Typically, the molecules will be 25 tested over a range of concentrations, and a series of control wells lacking one or more elements of the test assays can be used for accuracy in the evaluation of the results. An alternative to this method involves reversing the "positions" of the proteins, i.e., 30 immobilizing a B7-like binding partner to the microtiter plate wells, incubating with the test molecule and radiolabeled B7-like polypeptide,

20

25

30

A-692

determining the extent of B7-like polypeptide binding. See, for example, Chapter 18, Current Protocols in Molecular Biology, Ausubel et al., eds., John Wiley & Sons, New York, NY (1995).

5 As an alternative to radiolabelling, a B7-like polypeptide or its binding partner may be conjugated to biotin and the presence of biotinylated protein can then be detected using streptavidin linked enzyme, such horseradish peroxidase as (HRP) 10 alkaline phosphatase (AP), that can be detected colorometrically, or by fluorescent tagging streptavidin. An antibody directed to a B7-like polypeptide or to a B7-like binding partner conjugated to biotin may also be used and can be 15 detected after incubation with enzyme-linked streptavidin linked to AP or HRP.

A B7-like polypeptide or a B7-like binding partner can also be immobilized by attachment to agarose beads, acrylic beads or other types of such inert solid phase The substrate-protein complex can substrates. placed in a solution containing the complementary protein and the test compound. After incubation, the beads can be precipitated by centrifugation, and the amount of binding between a B7-like polypeptide and its binding partner can be assessed using the methods described herein. Alternatively, the substrate-protein complex can be immobilized in a column, and the test molecule and complementary protein are passed through The formation of a complex between a B7the column. like polypeptide and its binding partner can then be assessed using any of the techniques set forth herein, i.e., radiolabelling, antibody binding, or the like.

Another in vitro assay that is useful for identifying test molecule which a increases decreases the formation of a complex between a B7-like polypeptide and a B7-like binding partner is a surface plasmon resonance detector system such as the BIAcore assay system (Pharmacia, Piscataway, NJ). The BIAcore system may be carried out using the manufacturer's protocol. This assay essentially involves the covalent binding of either a B7-like polypeptide or a B7-like binding partner to a dextran-coated sensor chip which 10 is located in a detector. The test compound and the other complementary protein can then be injected, either simultaneously or sequentially, into the chamber containing the sensor chip. The amount complementary protein that binds can be assessed based 15 on the change in molecular mass which is physically associated with the dextran-coated side of the sensor chip; the change in molecular mass can be measured by the detector system.

In some cases, it may be desirable to evaluate two or more test compounds together for their ability to increase or decrease the formation of a complex between a B7-like polypeptide and a B7-like binding partner. In these cases, the assays set forth herein can be readily modified by adding such additional test compound(s) either simultaneous with, or subsequent to, the first test compound. The remainder of the steps in the assay are as set forth herein.

In vitro assays such as those described herein may
30 be used advantageously to screen large numbers of
compounds for effects on complex formation by a B7-like
polypeptide and B7-like binding partner. The assays

may be automated to screen compounds generated in phage display, synthetic peptide, and chemical synthesis libraries.

Compounds which increase or decrease the formation of a complex between a B7-like polypeptide and a B7-5 like binding partner may also be screened in cell culture using cells and cell lines expressing either a B7-like polypeptide or a B7-like binding partner. Cells and cell lines may be obtained from any mammal, 10 but preferably will be from human or other primate, canine, or rodent sources. The binding of a B7-like polypeptide to cells expressing B7-like binding partner at the surface is evaluated in the presence or absence of test molecules, and the extent of binding may be 15 determined by, for example, flow cytometry using a biotinylated antibody to a B7-like binding partner. Cell culture assays can be used advantageously further evaluate compounds that score positive protein binding assays described herein.

20 Cell cultures can also be used to screen the impact of drug candidate. a For example, candidates may decrease or increase the expression of a B7-like gene. In certain embodiments, the amount of B7-like polypeptide that is produced may be measured 25 after exposure of the cell culture to the candidate. In certain embodiments, one may detect the actual impact of the drug candidate on the cell For example, the overexpression particular gene may have a particular impact on the 30 cell culture. In such cases, one may test a drug candidate's ability to increase or decrease expression of the gene or its ability to prevent or

inhibit a particular impact on the cell culture. In other examples, the production of a particular metabolic product such as a fragment of a polypeptide, may result in, or be associated with, a disease or pathological condition. In such cases, one may test a drug candidate's ability to decrease the production of such a metabolic product in a cell culture.

Internalizing Proteins

The tat protein sequence (from HIV) can be used to internalize proteins into a cell. See e.g., Falwell et 10 al., Proc. Natl. Acad. Sci., 91:664-668 (1994). For example, an 11 amino acid sequence (YGRKKRRQRRR) of the tat protein (termed the "protein transduction domain", or TAT PDT) has been described as mediating 15 delivery across the cytoplasmic membrane and nuclear membrane of a cell. See Schwarze et al., Science, 285:1569-1572 (1999); and Nagahara et al., Nature Medicine, 4:1449-1452 (1998). In procedures, FITC-constructs (FITC-GGGGYGRKKRRQRRR) are 20 prepared which bind to cells as observed fluorescence-activated cell sorting (FACS) analysis, and these constructs penetrate tissues after adminstration. Next, tat-bgal fusion proteins are constructed. Cells treated with this construct demonstrated b-gal activity. Following injection, a 25 number of tissues, including liver, kidney, lung, heart, and brain tissue have been found to demonstrate expression using these procedures. It is believed that these constructions underwent some degree of unfolding in order to enter the cell; as such, refolding may be 30 required after entering the cell.

It will thus be appreciated that the tat protein

sequence may be used to internalize a desired protein or polypeptide into a cell. For example, using the tat protein sequence, a B7-like antagonist (such as anti-B7-like selective binding agent, small molecule, soluble receptor, or antisense oligonucleotide) can be administered intracellularly to inhibit the activity of the mature form of at least one B7-like molecule. used herein, the term "B7-like molecules" refers to B7-like nucleic acid molecules and both B7-like polypeptides as defined herein. Where desired, the B7like proteins may also be internally administered to a cell using these procedures. See also, Strauss, E., "Introducing Proteins Into the Body's Cells", Science, 285:1466-1467 (1999).

15

20

25

30

10

5

Therapeutic Uses

Polypeptides of the invention, and agonists and antagonists thereof, may be used to regulate T-cell function. Accordingly, B7-like polypeptides may be used to treat, diagnose, ameliorate, or prevent acute or chronic diseases associated with T-cell function.

Agonists and antagonists include those molecules which regulate B7-like polypeptide activity and either increase or decrease at least one activity of the mature form of at least one B7-like polypeptide such as one activity associated with T-cell functions. example, T-cell activation. Agonists or antagonists co-factors, be such as a protein, peptide, carbohydrate, lipid or small molecular weight molecule, which interact with at least one B7-like polypeptide and thereby regulate polypeptide activity. Potential polypeptide agonists or antagonists include antibodies

5

10

that react with either soluble or membrane-bound forms of at least one B7-like polypeptide which comprise part or all of the extracellular domains of the said proteins. Molecules that regulate B7-like polypeptide expression typically include nucleic acids that can act as anti-sense regulators of expression.

The phenotype of transgenic mice expressing B7-like polypeptide corresponding to SEQ ID NO: 14 showed seminal vesicle hyperplasia. Accordingly, agonists and antagonists of B7-like polypeptide activity may be useful in the treatment of reproductive disorders and proliferative disorders.

Antibodies, soluble proteins comprising for example extracellular domains, and other regulators of 15 B7-like polypeptide expression that result in prolonged or enhanced T-cell activation can be used to increase the immune response to tumors. B7-like polypeptides may play a role in the growth and maintenance of cancer cells based on overexpression causing seminal vesicle hyperplasia. Accordingly, agonists or antagonists to 20 B7-like polypeptides may be useful for the diagnosis and/or treatment of cancer. Examples of such cancers include, but are not limited to, seminal vesicle cancer, lung cancer, brain cancer, breast cancer, cancers of the hematopoetic system, prostate cancer, 25 ovarian cancer, and testicular cancer. Other cancers are encompassed within the scope of the invention. B7-like polypeptide pathway can also be manipulated to regulate CTL response in a number of other clinical settings, including allograft transplantation, graft 30 vs. host disease, and autoimmune diseases.

B7-like polypeptides may play a role in the inappropriate proliferation of cells based on overexpression causing seminal vesicle hyperplasia.

Accordingly, agonists or antagonists to B7-like polypeptides may be useful for the diagnosis and/or treatment of diseases where there is abnormal cell proliferation. Examples of such diseases include, but are not limited to, arteriosclerosis and vascular restenosis. Other diseases influenced by the inappropriate proliferation of cells are encompassed within the scope of the invention.

B7-like polypeptides may play a role in 10 reproductive system based on overexpression causing seminal vesicle hyperplasia. Accordingly, agonists or antagonists to B7-like polypeptides may be useful for of diagnosis and/or treatment reproductive disorders. Examples of such diseases include, but are 15 not limited to, infertility, miscarriage, preterm labor and delivery, and endometriosis. Other diseases of the reproductive system are encompassed within the scope of the invention.

B7-like polypeptides, and agonists and antagonists 20 thereof, may be used in the treatment of autoimmune disease, graft survival, immune cell activation for inhibiting tumor cell growth, T-cell dependent B-cell mediated diseases, and cancer gene immunotherapy. one embodiment, antagonists or inhibitors of B7-like 25 polypeptide function may be beneficial to alleviate chronic in diseases with symptoms immune dysfunction. Autoimmune diseases, such as systemic erythematosis, rheumatoid arthritis, thrombocytopenic purpura (ITP) and psoriasis may be 30 B7-like treated with polypeptide antagonists inhibitors. In addition, chronic inflammatory diseases, such as inflammatory bowel disease (Crohn's disease and ulcerative colitis), Grave's disease,

10

15

20

25

30

Hashimoto's thyroiditis and diabetes mellitus may also be treated with B7-like polypeptide inhibitors.

B7-like polypeptide antagonists may be used as immunosuppressive agents for bone marrow and organ transplantation and may be used to prolong graft survival. Such antagonists may provide significant advantages over existing treatment. Bone marrow and organ transplantation therapy must contend with T-cell mediated rejection of the foreign cells or tissue by the host. Present therapeutic regimens for inhibiting T-cell mediated rejection involve treatment with the cyclosporine or FK506. While drugs are effective, patients suffer from serious side effects, including hepatotoxicity, nephrotoxicity neurotoxicity. The target for the cyclosporin/FK506 class of therapeutics is calcineurin, a phosphatase with ubiquitous expression. Inhibitors of B7-like polypeptides or proteins may lack the severe side effects observed with use of the immunotherapeutic agents.

Antagonists of B7-like polypeptides or proteins may be used as immunosuppressive agents for autoimmune disorders, such as rheumatoid arthritis, psoriasis, multiple sclerosis, diabetes, and systemic lupus erythematosus.

Antagonists of the B7-like polypeptides or proteins may also be used to alleviate toxic shock syndrome, inflammatory bowel disease, allosensitization due to blood transfusions, T-cell dependent B-cell mediated diseases, and the treatment of graft vs. host disease.

Gene therapy using B7-like polypeptide or protein genes of the invention may be used in cancer

5

10

immunotherapy. B7-like polypeptide genes introduced into cancer cells can transform them into antigen presenting cells that can be recognized by the T-cells of the immune system when introduced back into animal. Recognition of the transfected tumor cells by the T-cells results in eradication of both tumors cells expressing, or not expressing, the B7-like polypeptide This immunotherapy approach may be used for various leukemias. sarcomas, melanomas, adenocarcinomas, breast carcinomas, prostate tumors, lung carcinomas, colon carcinomas and other tumors. This invention encompasses using the B7-like polypeptide gene in a similar manner to enhance T-cell activation in response to variety of tumors.

For instance, many vaccines act by eliciting an effective and specific antibody response. Some vaccines, especially those against intestinal microorganisms (e.g. Hepatitis A virus, and Salmonellas), elicit a short-lived antibody response. It is desirable to potentiate and prolong this response in order to increase the effectiveness of the vaccine. Therefore, soluble B7-like polypeptides or proteins may serve as a vaccine adjuvant.

Anti-viral responses may also be enhanced by activators or agonists of the B7-like protein pathway. The enhancement of cellular immune functions by B7-like polypeptide or protein/-Fc, may also be beneficial in eliminating virus-infected cells. In a complementary fashion, B7-like polypeptide or protein/-Fc may also have effects on humoral immune functions that may enhance antibody mediated responses and that may function to help clear free-virus from the body.

Conversely, there are a number of clinical conditions that would be ameliorated by the inhibition

5

10

15

20

25

30

of antibody production. Hypersensitivity is a normally beneficial immune response that is exaggerated or inappropriate, and leads to inflammatory reactions and tissue damage. Hypersensitivity reactions which are antibody-mediated may be particularly susceptible to antagonism by inhibitors of B7-like polypeptide or protein activity. Allergies, hay fever, asthma and acute edema cause type-I hypersensitivity reactions, and these reactions may be suppressed by protein, antibody or small molecule inhibitors of B7-like polypeptide or protein activity.

antibody-mediated Diseases that cause hypersensitivity reactions, including systemic lupus arthritis erythematosis, (rheumatoid arthritis, reactive arthritis, psoriatic arthritis), nephropathies (glomerulo-nephritis, membranous, mesangiocapillary, segmental, focal necrotizing, crescentic, proliferative tubulopathies), skin disorders (pemphigus and pemphigoid, erythema nodosum), endocrinopathies (Grave's disease, Hashimoto's thyroiditis and diabetes mellitus), various pneumopathies (especially extrinsic alveolitis), various vasculopathies, coeliac disease, with aberrant production of IgA, many anemias thrombocytopenias, Guillain-Barre syndrome, and myasthenia gravis may be treated with B7-like polypeptide or protein antagonists.

In addition, lymphoproliferative disorders, such as multiple myeloma, Waldenstrom's macroglobulinemia and crioglobulinemias may be inhibited by protein, antibody or small molecule antagonists of B7-like polypeptides or proteins.

Finally, graft versus host disease, an "artificial" immune disorder, may benefit from the

inhibition of antibody production by B7-like polypeptide or protein antagonists.

Other diseases associated with undesirable levels of one or more of the receptors of the present B7-like protein, and/or the present B7-like protein itself, are encompassed within the scope of the invention. Undesirable levels include excessive and/or sub-normal levels of the ligand of the present B7-like protein, and/or the B7-like protein described herein.

10 B7-like polypeptides, proteins, agonists and antagonists may be used in combination with cytokines, growth factors, antibiotics, anti-inflammatories, and/or chemotherapeutic agents as is appropriate for the indication being treated.

15

20

25

5

B7-like Compositions and Administration

Therapeutic compositions are within the scope of the present invention. Such B7-like pharmaceutical compositions may comprise a therapeutically effective amount of a B7-like polypeptide or a B7-like nucleic acid molecule in admixture with a pharmaceutically or physiologically acceptable formulation agent selected for suitability with the mode of administration. Pharmaceutical compositions may comprise therapeutically effective amount of one or more B7-like with agents in admixture selective binding physiologically acceptable pharmaceutically or formulation agent selected for suitability with the mode of administration.

Acceptable formulation materials preferably are nontoxic to recipients at the dosages and concentrations employed.

The pharmaceutical composition may contain 5 formulation materials for modifying, maintaining or preserving, for example, the pH, osmolarity, viscosity, clarity, color, isotonicity, odor, sterility, stability, rate of dissolution or release, adsorption penetration of the composition. Suitable 10 formulation materials include, but are not limited to, amino acids (such as glycine, glutamine, asparagine, arginine or lysine), antimicrobials, antioxidants (such as ascorbic acid, sodium sulfite or sodium hydrogensulfite), buffers (such as borate, bicarbonate, Tris-15 HCl, citrates, phosphates, other organic acids), bulking agents (such as mannitol or glycine), chelating agents (such as ethylenediamine tetraacetic acid (EDTA)), complexing agents (such as caffeine, polyvinylpyrrolidone, beta-cyclodextrin 20 hydroxypropyl-beta-cyclodextrin), fillers. monosaccharides, disaccharides, and other carbohydrates (such as glucose, mannose, or dextrins), proteins (such as serum albumin, gelatin or immunoglobulins), coloring, flavoring and diluting agents, emulsifying 25 agents, hydrophilic polymers (such polyvinylpyrrolidone), low molecular weight polypeptides, salt-forming counterions (such sodium), preservatives (such as benzalkonium chloride, benzoic acid, salicylic acid, thimerosal, phenethyl alcohol, methylparaben, propylparaben, chlorhexidine, 30 sorbic acid or hydrogen peroxide), solvents (such as glycerin, propylene glycol or polyethylene glycol), sugar alcohols (such mannitol or as

sorbitol),

10

suspending agents, surfactants or wetting agents (such as pluronics, PEG, sorbitan esters, polysorbates such polysorbate as 20, polysorbate 80, triton, tromethamine, lecithin, cholesterol, tyloxapal), stability enhancing agents (sucrose or sorbitol), tonicity enhancing agents (such as alkali metal halides (preferably sodium or potassium chloride), mannitol sorbitol), delivery vehicles, diluents, excipients and/or pharmaceutical adjuvants. (See Remington's Pharmaceutical Sciences, 18th Edition, A.R. Gennaro, ed., Mack Publishing Company (1990)).

The optimal pharmaceutical composition will be determined by one skilled in the art depending upon, for example, the intended route of administration, delivery format, and desired dosage. See for example, Remington's Pharmaceutical Sciences, supra. Such compositions may influence the physical state, stability, rate of in vivo release, and rate of in vivo clearance of the B7-like molecule.

20 The primary vehicle or carrier in a pharmaceutical composition may be either aqueous or non-aqueous in nature. For example, a suitable vehicle or carrier may be water for injection, physiological saline solution, artificial cerebrospinal fluid, possibly 25 supplemented with other materials common compositions for parenteral administration. buffered saline or saline mixed with serum albumin are exemplary vehicles. further Other exemplary pharmaceutical compositions comprise Tris buffer of about pH 7.0-8.5, or acetate buffer of about pH 4.0-30 5.5, which may further include sorbitol or a suitable substitute therefor. In one embodiment of the present

5

25

30

invention, B7-like polypeptide compositions may be prepared for storage by mixing the selected composition having the desired degree of purity with optional formulation agents (Remington's Pharmaceutical Sciences, supra) in the form of a lyophilized cake or an aqueous solution. Further, the B7-like polypeptide product may be formulated as a lyophilizate using appropriate excipients such as sucrose.

The B7-like pharmaceutical compositions can be selected for parenteral delivery. Alternatively, the compositions may be selected for inhalation or for delivery through the digestive tract, such as orally. The preparation of such pharmaceutically acceptable compositions is within the skill of the art.

The formulation components are present in concentrations that are acceptable to the site of administration. For example, buffers are used to maintain the composition at physiological pH or at slightly lower pH, typically within a pH range of from about 5 to about 8.

When parenteral administration is contemplated, the therapeutic compositions for use in this invention may be in the form of a pyrogen-free, parenterally acceptable aqueous solution comprising the desired B7-like molecule in a pharmaceutically acceptable vehicle. A particularly suitable vehicle for parenteral injection is sterile distilled water in which a B7-like molecule is formulated as a sterile, isotonic solution, properly preserved. Yet another preparation can involve the formulation of the desired molecule with an agent, such as injectable microspheres, bio-erodible particles, polymeric compounds (polylactic acid,

5

polyglycolic acid), or beads, or liposomes, that provides for the controlled or sustained release of the product which may then be delivered as a injection. Hyaluronic acid may also be used, and this may have the effect of promoting sustained duration in the circulation. Other suitable means for introduction of the desired molecule include implantable drug delivery devices.

In one embodiment, a pharmaceutical composition 10 may be formulated for inhalation. For example, a B7like molecule may be formulated as a dry powder for inhalation. B7-like polypeptide or B7-like nucleic molecule inhalation solutions acid may also formulated with a propellant for aerosol delivery. 15 yet another embodiment, solutions may be nebulized. Pulmonary administration is further described in PCT application no. PCT/US94/001875, which describes pulmonary delivery of chemically modified proteins.

It is also contemplated that certain formulations 20 may be administered orally. In one embodiment of the present invention, B7-like molecules which administered in this fashion can be formulated with or without those carriers customarily used compounding of solid dosage forms such as tablets and 25 capsules. For example, a capsule may be designed to release the active portion of the formulation at the point in the gastrointestinal tract bioavailability is maximized and pre-systemic degradation is minimized. Additional agents can be 30 included to facilitate absorption of the B7-like molecule. Diluents, flavorings, low melting point waxes, vegetable oils, lubricants, suspending agents, tablet disintegrating agents, and binders may also be

employed. Another pharmaceutical composition may involve an effective quantity of B7-like molecules in a mixture with non-toxic excipients which are suitable for the manufacture of tablets. By dissolving the tablets in sterile water, or other appropriate vehicle, 5 solutions can be prepared in unit dose form. Suitable excipients include, but are not limited to, inert diluents, such as calcium carbonate, sodium carbonate or bicarbonate, lactose, or calcium phosphate; binding agents, such as starch, gelatin, or acacia; or 10 lubricating agents such as magnesium stearate, stearic acid, talc. Additional or B7-like pharmaceutical compositions will be evident to those skilled in the art, including formulations involving B7-like 15 polypeptides in sustained- or controlled-delivery formulations. Techniques for formulating a variety of other sustained- or controlled-delivery means, such as liposome carriers, bio-erodible microparticles porous beads and depot injections, are also known to 20 those skilled in the art. See for example, PCT/US93/00829 which describes controlled release of porous polymeric microparticles for the delivery of pharmaceutical compositions. Additional examples of sustained-release preparations include semipermeable 25 polymer matrices in the form of shaped articles, e.g. films, or microcapsules. Sustained release matrices may include polyesters, hydrogels, polylactides (U.S. 3,773,919 and EP 58,481), copolymers of L-glutamic acid gamma ethyl-L-glutamate (Sidman et al., 30 Biopolymers, 22:547-556 (1983)), poly (2-hydroxyethylmethacrylate) (Langer et al., J. Biomed. Mater. Res., 15:167-277 (1981) and Langer, Chem. Tech., 12:98-105 (1982)), ethylene vinyl acetate (Langer et al., supra)

5

10

15

30

or poly-D(-)-3-hydroxybutyric acid (EP 133,988). Sustained-release compositions also may include liposomes, which can be prepared by any of several methods known in the art. See e.g., Eppstein et al., Proc. Natl. Acad. Sci. USA, 82:3688-3692 (1985); EP 36,676; EP 88,046; EP 143,949.

The B7-like pharmaceutical composition to be used for in vivo administration typically must be sterile. This may be accomplished by filtration through sterile filtration membranes. Where the composition lyophilized, sterilization using these methods may be conducted either prior to, or following, lyophilization and reconstitution. The composition for parenteral administration may be stored in lyophilized form or in solution. In addition, parenteral compositions generally are placed into a container having a sterile access port, for example, an intravenous solution bag or vial having a stopper pierceable by a hypodermic injection needle.

Once the pharmaceutical composition has been formulated, it may be stored in sterile vials as a solution, suspension, gel, emulsion, solid, or a dehydrated or lyophilized powder. Such formulations may be stored either in a ready-to-use form or in a form (e.g., lyophilized) requiring reconstitution prior to administration.

In a specific embodiment, the present invention is directed to kits for producing a single-dose administration unit. The kits may each contain both a first container having a dried protein and a second container having an aqueous formulation. Also included within the scope of this invention are kits containing

5

10

15

20

25

30

single and multi-chambered pre-filled syringes (e.g., liquid syringes and lyosyringes).

An effective amount of a B7-like pharmaceutical composition to be employed therapeutically will depend, for example, the therapeutic upon context objectives. One skilled in the art will appreciate that the appropriate dosage levels for treatment will thus vary depending, in part, upon the molecule delivered. the indication for which the molecule is being used, the route of administration, and the size (body weight, body surface or organ size) and condition (the age and general health) of the Accordingly, the clinician may titer patient. dosage and modify the route of administration to obtain the optimal therapeutic effect. A typical dosage may range from about 0.1 μ g/kg to up to about 100 mg/kg or more, depending on the factors mentioned above. other embodiments, the dosage may range from 0.1 $\mu g/kg$ to about 100 mg/kg; or 1 μ g/kg up 100 mg/kg; or 5 μ g/kg up to about 100 mg/kg.

The frequency of dosing will depend upon the pharmacokinetic parameters of the B7-like molecule in the formulation used. Typically, a clinician will administer the composition until a dosage is reached that achieves the desired effect. The composition may therefore be administered as a single dose, or as two or more doses (which may or may not contain the same amount of the desired molecule) over time, or as a continuous infusion via implantation device catheter. Further refinement of the appropriate dosage is routinely made by those of ordinary skill in the art and is within the ambit of tasks routinely performed by

5

10

15

20

25

30

them. Appropriate dosages may be ascertained through use of appropriate dose-response data.

The route of administration of the pharmaceutical composition is in accord with known methods, e.g. oral, injection by intravenous, intraperitoneal, intracerebral (intra-parenchymal), intracerebroventricular, intramuscular, intra-ocular, intraarterial, intraportal, or intralesional routes, or by sustained release systems or implantation device. Where desired, the compositions may be administered by bolus injection or continuously by infusion, or by implantation device.

Alternatively or additionally, the composition may be administered locally via implantation of a membrane, sponge, or other appropriate material on to which the desired molecule has been absorbed or encapsulated. Where an implantation device is used, the device may be implanted into any suitable tissue or organ, and delivery of the desired molecule may be via diffusion, timed release bolus, or continuous administration.

In some cases, it may be desirable to use B7-like pharmaceutical compositions in an ex vivo manner. In such instances, cells, tissues, or organs that have been removed from the patient are exposed to B7-like pharmaceutical compositions after which the cells, tissues and/or organs are subsequently implanted back into the patient.

In other cases, a B7-like polypeptide can be delivered by implanting certain cells that have been genetically engineered, using methods such as those described herein, to express and secrete the

5

10

15

20

polypeptide. Such cells may be animal or human cells, and may be autologous, heterologous, or xenogeneic. Optionally, the cells may be immortalized. In order to decrease the chance of an immunological response, the cells may be encapsulated to avoid infiltration of surrounding tissues. The encapsulation materials are typically biocompatible, semi-permeable polymeric enclosures or membranes that allow the release of the protein product(s) but prevent the destruction of the cells by the patient's immune system or by other detrimental factors from the surrounding tissues.

Additional embodiments of the present invention relate to cells and methods (e.g., homologous recombination and/or other recombinant production in vitro production methods) for both the therapeutic polypeptides and for the production and delivery of therapeutic polypeptides by gene therapy or Homologous and other recombination therapy. methods may be used to modify a cell that contains a normally transcriptionally silent B7-like gene, or an under-expressed gene, and thereby produce a cell which expresses therapeutically efficacious amounts of B7like polypeptides.

Homologous recombination is a technique originally developed for targeting genes to induce or correct mutations in transcriptionally active genes (Kucherlapati, Prog. in Nucl. Acid Res. & Mol. Biol., 36:301, 1989). The basic technique was developed as a method for introducing specific mutations into specific regions of the mammalian genome (Thomas et al., Cell, 44:419-428, 1986; Thomas and Capecchi, Cell, 51:503-512, 1987 and Doetschman et al., Proc. Natl. Acad.

30

A-692

Sci., 85:8583-8587, 1988) or to correct specific mutations within defective genes (Doetschman et al., Nature, 330:576-578, 1987). Exemplary homologous recombination techniques are described in U.S. Patent No. 5,272,071 (EP 9193051, EP Publication No. 505500; PCT/US90/07642, International Publication No. WO 91/09955).

Through homologous recombination, the DNA sequence to be inserted into the genome can be directed to a 10 specific region of the gene of interest by attaching it to targeting DNA. The targeting DNA is a nucleotide sequence that is complementary (homologous) to a region of the genomic DNA. Small pieces of targeting DNA that are complementary to a specific region of the genome are put in contact with the parental strand during the 15 DNA replication process. It is a general property of DNA that has been inserted into a cell to hybridize, and therefore, recombine with other pieces endogenous DNA through shared homologous regions. Τf 20 this complementary strand is attached an oligonucleotide that contains a mutation or a different an additional nucleotide, it sequence or incorporated into the newly synthesized strand as a result of the recombination. As a result of the proofreading function, it is possible for the 25 sequence of DNA to serve as the template. transferred DNA is incorporated into the genome.

Attached to these pieces of targeting DNA are regions of DNA which may interact with or control the expression of a B7-like polypeptide, e.g., flanking sequences. For example, a promoter/enhancer element, a suppresser, or an exogenous transcription modulatory

5

10

element is inserted in the genome of the intended host proximity and orientation sufficient cell in influence the transcription of DNA encoding the desired B7-like polypeptide. The control element controls a portion of the DNA present in the host cell genome. Thus, the expression of the desired B7-like polypeptide may be achieved not by transfection of DNA that encodes the B7-like gene itself, but rather by the use of targeting DNA (containing regions of homology with the endogenous gene of interest) coupled with regulatory segments that provide the endogenous gene sequence with recognizable signals for transcription of a B7-like polypeptide.

In exemplary method, the expression of desired targeted gene in a cell (i.e., a desired 15 endogenous cellular gene) is altered via homologous recombination into the cellular genome at a preselected site, by the introduction of DNA which includes at least a regulatory sequence, an exon and a splice donor 20 site. These components are introduced into chromosomal (genomic) DNA in such a manner that this, effect, in results in the production of a transcription unit (in which the regulatory sequence, the exon and the splice donor site present in the DNA construct are operatively linked to the endogenous 25 As a result of the introduction of these gene). components into the chromosomal DNA, the expression of the desired endogenous gene is altered.

Altered gene expression, as described herein,
30 encompasses activating (or causing to be expressed) a
gene which is normally silent (unexpressed) in the cell
as obtained, as well as increasing the expression of a

5

gene which is at physiologically not expressed significant levels in the cell as obtained. embodiments further encompass changing the pattern of regulation or induction such that it is different from the pattern of regulation or induction that occurs in cell as obtained, and reducing (including eliminating) the expression of а gene which expressed in the cell as obtained.

One method by which homologous recombination can 10 be used to increase, or cause, B7-like polypeptide production from a cell's endogenous B7-like gene involves first using homologous recombination to place recombination sequence from a site-specific recombination system (e.g., Cre/loxP, FLP/FRT) (Sauer, 15 Current Opinion In Biotechnology, 5:521-527, 1994 and Sauer, Methods InEnzymology, 225:890-900, upstream (that is, 5' to) of the cell's endogenous genomic B7-like polypeptide coding region. A plasmid containing a recombination site homologous to the site 20 that was placed just upstream of the genomic B7-like polypeptide coding region is introduced into modified cell line along with the appropriate recombinase enzyme. This recombinase causes the plasmid to integrate, via the plasmid's recombination 25 site, into the recombination site located just upstream of the genomic B7-like polypeptide coding region in the cell line (Baubonis and Sauer, Nucleic Acids Res., 21:2025-2029, 1993 and O'Gorman et al., Science, 251:1351-1355, 1991). Any flanking sequences known to 30 increase transcription (e.g., enhancer/promoter, intron, translational enhancer), if properly positioned in this plasmid, would integrate in such a manner as to

20

25

30

A-692

create a new or modified transcriptional unit resulting in *de novo* or increased B7-like polypeptide production from the cell's endogenous B7-like gene.

A further method to use the cell line in which the site specific recombination sequence had been placed 5 just upstream of the cell's endogenous genomic B7-like polypeptide coding region is to use homologous recombination to introduce a second recombination site elsewhere in the cell line's genome. The appropriate 10 recombinase enzyme is then introduced into the tworecombination-site cell line, causing a recombination event (deletion, inversion, translocation) Current Opinion In Biotechnology, supra, 1994; Sauer, Methods In Enzymology, supra, 1993) that would create a 15 new or modified transcriptional unit resulting in de novo or increased B7-like polypeptide production from the cell's endogenous B7-like gene.

An additional approach for increasing, or causing, the expression of B7-like polypeptide from a cell's endogenous B7-like gene involves increasing, causing, the expression of a gene or genes (e.g., transcription factors) and/or decreasing the expression of a gene or genes (e.g., transcriptional repressors) in a manner which results in de novo or increased B7like polypeptide production from the cell's endogenous B7-like gene. This method includes the introduction of non-naturally occurring polypeptide (e.g., polypeptide comprising a site specific DNA binding domain fused to a transcriptional factor domain) into the cell such that de novo or increased B7-like polypeptide production from the cell's endogenous B7like gene results.

25

30

A-692

The present invention further relates to DNA constructs useful in the method of altering expression target gene. In certain embodiments, exemplary DNA constructs comprise: (a) one or more targeting sequences; (b) a regulatory sequence; (c) an 5 exon; and (d) an unpaired splice-donor site. targeting sequence in the DNA construct directs the integration of elements (a)-(d) into a target gene in a cell such that the elements (b)-(d) are operatively linked to sequences of the endogenous target gene. 10 another embodiment, the DNA constructs comprise: one or more targeting sequences, (b) a regulatory sequence, (c) an exon, (d) a splice-donor site, (e) an intron, and (f) a splice-acceptor site, wherein the 15 targeting sequence directs the integration of elements (a) - (f)such that the elements of (b) - (f)are operatively linked to the endogenous gene. targeting sequence is homologous to the preselected in the cellular chromosomal DNA with 20 homologous recombination is to occur. In the construct, the exon is generally 3' of the regulatory sequence and the splice-donor site is 3' of the exon.

If the sequence of a particular gene is known, such as the nucleic acid sequence of B7-like polypeptide presented herein, a piece of DNA that is complementary to a selected region of the gene can be synthesized or otherwise obtained, such by appropriate restriction of the native DNA at specific recognition sites bounding the region of interest. This piece serves as a targeting sequence(s) insertion into the cell and will hybridize to homologous region within the genome. Ιf this hybridization occurs during DNA replication, this piece

5

10

15

20

25

of DNA, and any additional sequence attached thereto, will act as an Okazaki fragment and will be incorporated into the newly synthesized daughter strand of DNA. The present invention, therefore, includes nucleotides encoding a B7-like polypeptide, which nucleotides may be used as targeting sequences.

B7-like polypeptide cell therapy, e.g., the implantation of cells producing B7-like polypeptides, also contemplated. This embodiment involves implanting cells capable of synthesizing and secreting a biologically active form of B7-like polypeptide. Such B7-like polypeptide-producing cells can be cells that are natural producers of B7-like polypeptides or may be recombinant cells whose ability to produce B7like polypeptides has been augmented by transformation with a gene encoding the desired B7-like polypeptide or with a gene augmenting the expression of B7-like polypeptide. Such a modification may be accomplished by means of a vector suitable for delivering the gene as well as promoting its expression and secretion. order to minimize a potential immunological reaction in patients being administered a B7-like polypeptide, as may occur with the administration of a polypeptide of a foreign species, it is preferred that the natural cells producing B7-like polypeptide be of human origin and produce human B7-like polypeptide. Likewise, it is preferred that the recombinant cells producing B7-like polypeptide be transformed with an expression vector containing a gene encoding a human B7-like polypeptide.

Implanted cells may be encapsulated to avoid the infiltration of surrounding tissue. Human or non-human animal cells may be implanted in patients in

biocompatible, semipermeable polymeric enclosures or membranes that allow the release of B7-like polypeptide, but that prevent the destruction of the cells by the patient's immune system or by other detrimental factors from the surrounding Alternatively, the patient's own cells, transformed to produce B7-like polypeptides ex vivo, may be implanted directly into the patient without such encapsulation.

Techniques for the encapsulation of living cells 10 are known in the art, and the preparation of the encapsulated cells and their implantation in patients may be routinely accomplished. For example, Baetge et al. (WO95/05452; PCT/US94/09299) describe membrane capsules containing genetically engineered cells for 15 biologically the effective delivery of active molecules. The capsules are biocompatible and are easily retrievable. The capsules encapsulate cells transfected with recombinant DNA molecules comprising DNA sequences coding for biologically active molecules 20 operatively linked to promoters that are not subject to down regulation in vivo upon implantation into a mammalian host. The devices provide for the delivery of the molecules from living cells to specific sites within a recipient. In addition, see U.S. Patent Nos. 4,892,538, 5,011,472, and 5,106,627. 25 A system for encapsulating living cells is described in PCT Application no. PCT/US91/00157 (Aebischer et al). See also, PCT Application no. PCT/US91/00155 Aebischer et al., Winn et al., Exper. Neurol., 113:322-329 (1991), 30 Aebischer et al., Exper. Neurol., 111:269-275 (1991);

and Tresco et al., ASAIO, 38:17-23 (1992).

15

25

30

A-692

In vivo and in vitro gene therapy delivery of B7like polypeptides is also envisioned. One example of a gene therapy technique is to use the B7-like gene genomic DNA, cDNA, and/or synthetic 5 encoding a B7-like polypeptide which may be operably linked to a constitutive or inducible promoter to form a "gene therapy DNA construct". The promoter may be homologous or heterologous to the endogenous B7-like gene, provided that it is active in the cell or tissue 10 type into which the construct will be inserted. Other components of the gene therapy DNA construct may optionally include, DNA molecules designed for sitespecific integration (e.g., endogenous sequences useful homologous recombination), tissue-specific promoter, enhancer(s) or silencer(s), DNA molecules capable of providing a selective advantage over the parent cell, DNA molecules useful as labels to identify transformed cells, negative selection systems, cell specific binding agents (as, for example, for cell 20 targeting), cell-specific internalization factors, and transcription factors to enhance expression by a vector as well as factors to enable vector manufacture.

Α gene therapy DNA construct can then be introduced into cells (either ex vivo or in vivo) using viral or non-viral vectors. One means for introducing the gene therapy DNA construct is by means of viral vectors as described herein. Certain vectors, such as retroviral vectors, will deliver the DNA construct to the chromosomal DNA of the cells, and the gene can integrate into the chromosomal DNA. Other vectors will function as episomes, and the gene therapy DNA construct will remain in the cytoplasm.

5

10

15

20

25

In yet other embodiments, regulatory elements can be included for the controlled expression of the B7like gene in the target cell. Such elements are turned on in response to an appropriate effector. In this way, a therapeutic polypeptide can be expressed when desired. One conventional control means involves the of small molecule dimerizers or rapalogs described in WO9641865 (PCT/US96/099486); WO9731898 (PCT/US97/03137) and WO9731899 (PCT/US95/03157)) used to dimerize chimeric proteins which contain a small molecule-binding domain and a domain capable initiating biological process, such as a DNA-binding protein or transcriptional activation protein. dimerization of the proteins can be used to initiate transcription of the transgene.

An alternative regulation technology uses a method of storing proteins expressed from the gene of interest inside the cell as an aggregate or cluster. interest is expressed as a fusion protein that includes a conditional aggregation domain which results in the retention of the aggregated protein in the endoplasmic reticulum. The stored proteins are stable and inactive inside the cell. The proteins can be released, however, by administering a drug (e.g., small molecule ligand) that conditional removes the aggregation domain and thereby specifically breaks apart the aggregates or clusters so that the proteins may be secreted from the cell. See, Science 287:816-817, and 826-830 (2000).

Other suitable control means or gene switches include, but are not limited to, the following systems.

Mifepristone (RU486) is used as a progesterone

5

antagonist. The binding of a modified progesterone receptor ligand-binding domain to the progesterone antagonist activates transcription by forming a dimer of two transcription factors which then pass into the nucleus to bind DNA. The ligand binding domain is modified to eliminate the ability of the receptor to bind to the natural ligand. The modified steroid hormone receptor system is further described in U.S. 5,364,791; WO9640911, and WO9710337.

- 10 Yet another control system uses ecdysone (a fruit fly steroid hormone) which binds to and activates an The receptor ecdysone receptor (cytoplasmic receptor). then translocates to the nucleus to bind a specific DNA response element (promoter from ecdysone-responsive 15 gene). The ecdysone receptor includes transactivation domain/DNA-binding domain/ligandbinding domain to initiate transcription. The ecdysone further described U.S. system is in 5,514,578; WO9738117; WO9637609; and WO9303162.
- 20 Another control means uses а positive tetracycline-controllable transactivator. This system involves a mutated tet repressor protein DNA-binding domain (mutated tet R-4 amino acid changes resulted in а reverse tetracycline-regulated 25 transactivator i.e., protein, it binds to a operator in the presence of tetracycline) linked to a polypeptide which activates transcription. systems are described in U.S. Patent Nos. 5,464,758; 5,650,298 and 5,654,168.
- Additional expression control systems and nucleic acid constructs are described in U.S. Patent Nos.

5

10

15

20

25

30

5,741,679 and 5,834,186, to Innovir Laboratories Inc.

In vivo gene therapy may be accomplished by introducing the gene encoding a B7-like polypeptide into cells via local injection of a B7-like nucleic acid molecule or by other appropriate viral or nonviral delivery vectors. See Hefti, Neurobiology, 25:1418-1435 (1994). For example, a nucleic acid molecule encoding a B7-like polypeptide contained in an adeno-associated virus (AAV) vector for delivery to the targeted cells (e.g., Johnson, International Publication No. W095/34670; International Application No. PCT/US95/07178). The recombinant AAV genome typically contains AAV inverted terminal repeats flanking a DNA sequence encoding a B7-like polypeptide operably linked to functional promoter and polyadenylation sequences.

Alternative suitable viral vectors include, are not limited to, retrovirus, adenovirus, herpes simplex virus, lentivirus, hepatitis virus, parvovirus, papovavirus, poxvirus, alphavirus, coronavirus, rhabdovirus, paramyxovirus, and papilloma vectors. U.S. Patent No. 5,672,344 describes an in vivo viral-mediated gene transfer system involving a recombinant neurotrophic HSV-1 vector. U.S. Patent No. 5,399,346 provides examples of a process for providing a patient with a therapeutic protein by the delivery of human cells which have been treated in vitro to insert DNA segment encoding а therapeutic protein. Additional methods and materials for the practice of gene therapy techniques are described in U.S. Patent No. 5,631,236 involving adenoviral vectors; U.S. Patent No. 5,672,510 involving retroviral vectors; and U.S.

5,635,399 involving retroviral vectors expressing cytokines.

Nonviral delivery methods include, but are not liposome-mediated transfer, limited to, naked DNA 5 delivery (direct injection), receptor-mediated transfer electroporation, (ligand-DNA complex), calcium phosphate precipitation, and microparticle bombardment (e.g., gene gun). Gene therapy materials and methods may also include the use of inducible promoters, 10 tissue-specific enhancer-promoters, DNA designed for site-specific integration, DNA sequences capable of providing a selective advantage over the parent cell, labels to identify transformed cells, negative selection systems and expression control 15 systems (safety measures), cell-specific binding agents (for cell targeting), cell-specific internalization transcription factors, and factors to enhance expression by a vector as well as methods of vector manufacture. Such additional methods and materials for 20 the practice of gene therapy techniques are described in U.S. Patent No. 4,970,154 involving electroporation techniques; W096/40958 involving nuclear ligands; U.S. 5,679,559 Patent No. describing lipoproteina containing system for gene delivery; U.S. Patent No. 25 5,676,954 involving liposome carriers; U.S. Patent No. 5,593,875 concerning methods for calcium phosphate transfection; and U.S. Patent No. 4,945,050 wherein biologically active particles are propelled at cells at a speed whereby the particles penetrate the surface of 30 the cells and become incorporated into the interior of the cells.

It is also contemplated that B7-like gene therapy

5

10

15

20

25

30

or cell therapy can further include the delivery of one or more additional polypeptide(s) in the same or a different cell(s). For example, the host cell may be modified to express and release both B7-like polypeptide and cytokines, growth factors and antiinflammatories. Alternatively, the B7-like polypeptide and cytokines, growth factors and anti-inflammatories, or B7-like polypeptide and cytokines, growth factors and anti-inflammatories, may be expressed in and released from separate cells.

Such cells may be separately introduced into the patient, or the cells may be contained in a single implantable device, such as the encapsulating membrane described above, or the cells may be separately modified by means of viral vectors.

A means to increase endogenous B7-like polypeptide expression in a cell via gene therapy is to insert one or more enhancer elements into the B7-like polypeptide promoter, where the enhancer element(s) can serve to increase transcriptional activity of the B7-like gene. The enhancer element(s) used will be selected based on the tissue in which one desires to activate the gene(s); enhancer elements known to confer promoter activation in that tissue will be selected. For example, if a gene encoding a B7-like polypeptide is to be "turned on" in T-cells, the 1ck promoter enhancer element may be used. Here, the functional portion of the transcriptional element to be added may be inserted fragment of DNA containing the polypeptide promoter (and optionally, inserted into a vector and/or 5' and/or 3' flanking sequence(s), etc.) using standard cloning techniques. This construct,

5

10

15

20

25

30

known as a "homologous recombination construct", can then be introduced into the desired cells either ex vivo or in vivo.

Gene therapy also can be used to decrease B7-like polypeptide expression by modifying the nucleotide sequence of the endogenous promoter(s). modification is typically accomplished via homologous recombination methods. For example, a DNA molecule containing all or a portion of the promoter of the B7for inactivation can like gene(s) selected be engineered to remove and/or replace pieces promoter that regulate transcription. For example the TATA box and/or the binding site of a transcriptional activator of the promoter may be deleted using standard molecular biology techniques; such deletion can inhibit promoter activity thereby repressing the transcription of the corresponding B7-like gene. The deletion of the TATA box or the transcription activator binding site in the promoter may be accomplished by generating a DNA construct comprising all or the relevant portion of the B7-like polypeptide promoter(s) (from the same or a related species as the B7-like gene(s) to be regulated) one or more of the TATAbox transcriptional activator binding site nucleotides are mutated via substitution, deletion and/or insertion of one or more nucleotides. As a result, the TATA box and/or activator binding site has decreased activity or is rendered completely inactive. The construct will typically contain at least about 500 bases of DNA that correspond to the native (endogenous) 5' and 3' DNA sequences adjacent to the promoter segment that has been modified. The construct may be introduced into the appropriate cells (either ex vivo or in vivo)

5

15

20

25

30

either directly or via a viral vector as described Typically, the integration of the construct herein. genomic DNA of the into the cells will via homologous recombination, where the 5' and 3′ DNA sequences in the promoter construct can serve to help modified integrate the promoter region via hybridization to the endogenous chromosomal DNA.

Additional Uses of B7-like Nucleic Acids and Polypeptides

Nucleic acid molecules of the present invention (including those that do not themselves encode biologically active polypeptides) may be used to map the locations of the B7-like gene and related genes on chromosomes. Mapping may be done by techniques known in the art, such as PCR amplification and in situ hybridization.

The B7-like polypeptides may be used (simultaneously or sequentially) in combination with one or more cytokines, growth factors, antibiotics, anti-inflammatories, and/or chemotherapeutic agents as is appropriate for the indication being treated.

Other methods may also be employed where it is desirable to inhibit the activity of one or more B7-like polypeptides. Such inhibition may be effected by nucleic acid molecules which are complementary to and hybridize to expression control sequences (triple helix formation) or to B7-like mRNA. For example, antisense DNA or RNA molecules, which have a sequence that is complementary to at least a portion of the selected B7-like gene(s) can be introduced into the cell. Antisense probes may be designed by available techniques

5

20

25

30

using the sequence of B7-like polypeptide disclosed herein. Typically, each such antisense molecule will be complementary to the start site (5' end) of each selected B7-like gene. When the antisense molecule then hybridizes to the corresponding B7-like mRNA, translation of this mRNA is prevented or reduced. Antisense inhibitors provide information relating to the decrease or absence of a B7-like polypeptide in a cell or organism.

10 Alternatively, gene therapy may be employed to create a dominant-negative inhibitor of one or more B7like polypeptides. In this situation, the DNA encoding mutant polypeptide of each selected B7-like polypeptide can be prepared and introduced into the cells of a patient using either viral or non-viral 15 methods as described herein. Each such mutant is typically designed to compete with endogenous polypeptide in its biological role.

addition, a B7-like polypeptide, In whether not, biologically active or may be used as immunogen, that is, the polypeptide contains at least to which antibodies be epitope may raised. one Selective binding agents that bind to B7-like а polypeptide (as described herein) may be used for in vivo and in vitro diagnostic purposes, including, but not limited to, use in labeled form to detect the presence of B7-like polypeptide in a body fluid or cell The antibodies may also be used to prevent, treat, or diagnose a number of diseases and disorders, including those recited herein. The antibodies may bind to a B7-like polypeptide so as to diminish or block at least one activity characteristic of a B7-like

polypeptide, or may bind to a polypeptide to increase at least one activity characteristic of a B7-like polypeptide (including by increasing the pharmacokinetics of the B7-like polypeptide).

While the present invention has been described in terms of the preferred embodiments, it is understood that variations and modifications will occur to those skilled in the art. Therefore, it is intended that the appended claims cover all such equivalent variations which come within the scope of the invention as claimed.